arXiv:1111.6100v2 [math.RA] 29 May 2012

THE DIXMIER CONJECTURE AND THE SHAPE OF POSSIBLE COUNTEREXAMPLES

JORGE A. GUCCIONE, JUAN J. GUCCIONE, AND CHRISTIAN VALQUI

Abstract. We introduce the notion of an irreducible pair of elements in the first Weyl algebra W. Then we show that each endomorphism of W which is not an automorphism, can be transformed into one that applies the canonical generators of W onto an irreducible pair with a specific shape (subrectangular). Consequently, the Dixmier conjecture is true if and only if there are no such irreducible pairs.

Introduction

The *Weyl algebra* A_1 over a field K is the quotient of the free associative and unital K-algebra on two generators X, Y by the ideal generated by the relation $[Y, X] = 1$. The Weyl algebra is the first of an infinite family of algebras, known as *Weyl algebras*, which were introduced by Hermann Weyl to study the Heisenberg uncertainty principle in quantum mechanics. The *n-th Weyl algebra over K* is the associative and unital K-algebra A_n generated by the $2n$ variables $X_1, Y_1, \dots, X_n, Y_n$, subject to the relations $[X_i, X_j] = [Y_i, Y_j] = 0$ and $[Y_i, X_j] = \delta_{ij}$, where δ_{ij} is the Krönecker symbol.

In [8] Dixmier posed six problems, the first of which, also known as the Dixmier conjecture, was the following: is an algebra endomorphism of the Weyl algebra A_1 on a characteristic zero field, necessarily an automorphism? This question makes sense for all the Weyl algebras, and in fact the generalized Dixmier conjecture claims that if $\mathrm{char}(K) = 0$, then any endomorphism of A_n must be an automorphism, for each $n \in \mathbb{N}$.

Currently, the Dixmier conjecture remains open even for the case $n = 1$. Some of the results in this topic are the following: In the early eighties, L. Vaserstein and V. Kac showed that the Dixmier conjecture implies the Jacobian conjecture (see [3], where this result is explicitly established). In 2005 the stable equivalence between the Dixmier and Jacobian conjectures was established by Yoshifumi Tsuchimoto [14], and the same result was obtained in an independent way in 2007 by Alexei Belov-Kanel and Maxim Kontsevich [6] and by Pascal Kossivi Adjamagbo and Arno van den Essen [2]. For a short proof of the equivalence between these two problems see [4]. In fact, in the [2] paper, there is also established the equivalence between the Dixmier conjecture and a similar conjecture about the endomorphisms of a family of Poisson algebras, that the autors call the Poisson Conjecture. Some other papers concerning the Dixmier conjecture in a direct or an indirect way are [5], [7] and [16].

In this work we deal with the case $n = 1$. From now on we write W (by Weyl) instead of A_1. A pair (P, Q) of elements of W is said to be *irreducible* if it is the

---

2010 *Mathematics Subject Classification.* Primary 16S32; Secondary 16W20.

Key words and phrases. Weyl algebra, Dixmier Conjecture.

Jorge A. Guccione and Juan J. Guccione research were supported by PIP 112-200801-00900 (CONICET).

Christian Valqui research was supported by PUCP-DGI-2011-0206 and Lucet 90-DAI-L005.

image of the pair (X,Y) of the canonical generators of W via an endomorphism which is not an automorphism, and it cannot be made "smaller" (in a sense that is established in detail in Definition 3.1) via automorphisms. Following the strategy of describing the generators of possible counterexamples, we prove the following result (Corollary 5.12): If the Dixmier conjecture is false, then there exist and irreducible pair (P,Q) such that the support of both P and Q is subrectangular (see Definition 1.5). This shape is the same achieved in the case of the Jacobian Conjecture (see for example [15, Corollary 10.2.21]).

In [8], the author uses in some proofs the geometric properties of the support of certain elements. In this paper we exploit such geometric properties. For each $(\rho,\sigma)\in\mathbb{Z}^2$ such that $\gcd(\rho,\sigma)=1$ and $\rho+\sigma\geq 0$ we consider the (ρ,σ)-valuation $v_{\rho,\sigma}$ introduced in [8]. One of the ingredients used by us is the concept of "leading term" $\ell_{\rho,\sigma}(P)$ of an element P of W with respect to the (ρ,σ)-valuation, which in [8] is called the (ρ,σ)-polynomial associated with P. In Proposition 1.9 we reprove some of the multiplicative properties of $\ell_{\rho,\sigma}(P)$, which are found partially in [8, Lemma 2.4]. Furthermore, in Definition 1.15, we introduce the notion of (ρ,σ)-bracket of a pair (P,Q) of elements of W, which, when it is non zero, coincides with the (ρ,σ)-polynomial associated with the commutator $[P,Q]$.

Our basic idea is to associate a polynomial $f_{P,\rho,\sigma}$ in one variable with $\ell_{\rho,\sigma}(P)$. On one hand this permits us to reprove some results of [8, Lemma 2.7] in Theorem 1.22 and, on the other hand, to write the (ρ,σ)-bracket $[P,Q]_{\rho,\sigma}$ as an expression of f_P, f_Q and its derivatives. This allows us to show that the (ρ,σ)-leading terms of the components of an irreducible pair (P,Q) must satisfy a very restrictive condition (Theorem 4.5): All irreducible factors of $f_{P,\rho,\sigma}$ and $f_{Q,\rho,\sigma}$ are factors of the polynomial $f_{F,\rho,\sigma}$, associated with a (ρ,σ)-homogeneous element $F\in W$ with (ρ,σ)-valuation $\rho+\sigma$. Then, using elementary automorphisms we can "cut" the form of the support of the components of an irreducible pair until they have the desired subrectangular shape. In the last section we "cut" the support further, and this allow us to prove that

$$B := \min\{\gcd(v_{1,1}(P), v_{1,1}(Q)),\ \text{where } (P,Q) \text{ is an irreducible pair}\}$$

is greater or equal to 9. The equivalent result for Jacobian pairs was proved by Nagata in [12], [13], based on a work of Appelgate and Onishi ([1]).

In general, the results about the shape in the commutative case and our case are very similar. We will compare the constructions described in [15] with ours. In the commutative case $\ell_{\rho,\sigma}(P)$ is called P_+ and the bracket corresponds to the Jacobian of P_+,Q_+. This is a bit simpler than our notation, but ignores the relation between the different valuations (called directions in [15]). Our Theorems 4.3 and 4.5 give an improved version of the corresponding result for the Jacobian conjecture, established in [15, Proposition 10.2.8]. On the commutative case, we also have the idea of a polynomial in one variable, for example in [15, Lemma 10.2.12], although this idea is not used in the proof of [15, Proposition 10.2.8], but is used in the proof of [15, Corollary 10.2.21]. The differential equation for the polynomials in item (1) of our Theorem 1.22 is also found partially in the definition of $\rho(z)$ before Lemma 10.2.13 in [15, page 250], but its importance is not fully noted.

In a forthcoming paper we will cut the subrectangular support further, which finally will leave very few possible shapes of irreducible pairs. In particular, we will improve the lower bound for the size of the support, and show a way to find possible counterexamples. If the same machinery works in the commutative case, it will improve substantively the lower bound for the degree of possible counterexamples established in [11], where it is shown that the degrees of the components P and Q of a Jacobian pair must be greater than 100.

1 Preliminaries

In this paper K is a characteristic zero field, W is the Weyl algebra on K, that is the unitary associative K-algebra generate by elements X, Y and the relation $[Y,X]=1$. Let $L := K[x,y]$ be the polynomial K-algebra in two variables and let $\Psi\colon W \to L$ be the K-linear map defined by $\Psi\bigl(X^iY^j\bigr) := x^iy^j$. Let

$$\overline{\mathfrak{V}} := \{(\rho,\sigma)\in\mathbb{Z}^2 : \gcd(\rho,\sigma)=1 \text{ and } \rho+\sigma\ge 0\}$$

and

$$\mathfrak{V} := \{(\rho,\sigma)\in\overline{\mathfrak{V}} : \rho+\sigma>0\}.$$

Note that $\overline{\mathfrak{V}} = \mathfrak{V}\cup\{(1,-1),(-1,1)\}$.

Definition 1.1. For all $(\rho,\sigma)\in\overline{\mathfrak{V}}$ and $(i,j)\in\mathbb{Z}\times\mathbb{Z}$ we write

$$v_{\rho,\sigma}(i,j) := \rho i + \sigma j.$$

Notations 1.2. Let $(\rho,\sigma)\in\overline{\mathfrak{V}}$. For $P=\sum a_{ij}x^iy^j \in L\setminus\{0\}$, we define:

- The *support* of P as
$$\operatorname{Supp}(P) := \{(i,j) : a_{ij}\ne 0\}.$$
- The *(ρ,σ)-degree* of P as $v_{\rho,\sigma}(P) := \max\{v_{\rho,\sigma}(i,j) : a_{ij}\ne 0\}$.
- The *(ρ,σ)-leading term* of P as
$$\ell_{\rho,\sigma}(P) := \sum_{\{\rho i+\sigma j = v_{\rho,\sigma}(P)\}} a_{ij}x^iy^j.$$
- $w(P) := (i_0, i_0 - v_{1,-1}(P))$ such that
$$i_0 = \max\{i : (i, i-v_{1,-1}(P)) \in \operatorname{Supp}(\ell_{1,-1}(P))\},$$
- $\ell_c(P) := a_{i_0j_0}$, where $(i_0,j_0)=w(P)$.
- $\ell_t(P) := a_{i_0j_0}x^{i_0}y^{j_0}$, where $(i_0,j_0)=w(P)$.
- $\overline{w}(P) := (i_0 - v_{-1,1}(P), i_0)$ such that
$$i_0 = \max\{i : (i-v_{-1,1}(P), i) \in \operatorname{Supp}(\ell_{-1,1}(P))\},$$
- $\overline{\ell}_c(P) := a_{i_0j_0}$, where $(i_0,j_0)=\overline{w}(P)$.
- $\overline{\ell}_t(P) := a_{i_0j_0}x^{i_0}y^{j_0}$, where $(i_0,j_0)=\overline{w}(P)$.

Notations 1.3. Let $(\rho,\sigma)\in\overline{\mathfrak{V}}$. For $P\in W\setminus\{0\}$, we define:

- The *support* of P as $\operatorname{Supp}(P) := \operatorname{Supp}\bigl(\Psi(P)\bigr)$.
- The *(ρ,σ)-degree* of P as $v_{\rho,\sigma}(P) := v_{\rho,\sigma}\bigl(\Psi(P)\bigr)$.
- The *(ρ,σ)-leading term* of P as $\ell_{\rho,\sigma}(P) := \ell_{\rho,\sigma}\bigl(\Psi(P)\bigr)$.
- $w(P) := w\bigl(\Psi(P)\bigr)$.
- $\ell_c(P) := \ell_c\bigl(\Psi(P)\bigr)$.
- $\ell_t(P) := \ell_c(P)X^{i_0}Y^{j_0}$, where $(i_0,j_0)=w(P)$.
- $\overline{w}(P) := \overline{w}\bigl(\Psi(P)\bigr)$.
- $\overline{\ell}_c(P) := \overline{\ell}_c\bigl(\Psi(P)\bigr)$.
- $\overline{\ell}_t(P) := \overline{\ell}_c(P)X^{i_0}Y^{j_0}$, where $(i_0,j_0)=\overline{w}(P)$.

Notation 1.4. We say that $P\in L$ is *(ρ,σ)-homogeneous* if $P=0$ or $P=\ell_{\rho,\sigma}(P)$.

Definition 1.5. We say that $P \in W$ is *subrectangular* if its support is contained in a rectangle who's farthest edge belongs to $\operatorname{Supp}(P)$. This means that there is $(i_0, j_0) \in \operatorname{Supp}(P)$ such that each $(i, j) \in \operatorname{Supp}(P)$ fulfills $i \le i_0$ and $j \le j_0$. In other words that $(i_0, j_0) \in \operatorname{Supp}(P)$ and that the support of P is contained in the rectangle whose vertices are $(0, 0)$, $(0, j_0)$, $(i_0, 0)$ and (i_0, j_0). Furthermore we will say that $P \in W$ is *subsquare* if $i_0 = j_0$.

Definition 1.6. Let $(\rho, \sigma) \in \overline{\mathfrak{V}}$ and let $P \in W \setminus \{0\}$.

- If $(\rho, \sigma) \neq (1, -1)$, then the *starting point of* P *with respect to* (ρ, σ) is
$$\operatorname{st}_{\rho,\sigma}(P) = w(\ell_{\rho,\sigma}(P)).$$
- If $(\rho, \sigma) \neq (-1, 1)$, then the *end point of* P *with respect to* (ρ, σ) is
$$\operatorname{en}_{\rho,\sigma}(P) = \overline{w}(\ell_{\rho,\sigma}(P)).$$

Note that $\operatorname{en}_{1,-1}(P) = w(P)$ and $\operatorname{st}_{-1,1}(P) = \overline{w}(P)$.



Illustration of some concepts introduced in Notation 1.3 and Definition 1.6 for
$$P = X^3 + X^5 + X^6Y + XY^3 + X^6Y^3 + X^3Y^4 + XY^6 + X^4Y^6 + X^2Y^7$$
and $(\rho, \sigma) = (3, 2)$. In this example
$$\ell_{\rho,\sigma}(P) = x^6y^3 + x^4y^6.$$

FIGURE 1

Lemma 1.7. *It is true that*
$$Y^jX^i = \sum_{k=0}^{j} k! \binom{j}{k} \binom{i}{k} X^{i-k}Y^{j-k}.$$

Proof. It follows easily using that
$$[Y, X^i] = iX^{i-1}, \qquad [Y^j, X^i] = [Y, X^i]Y^{j-1} + Y[Y^{j-1}, X^i]$$
and an induction argument. □

For $j \in \mathbb{Z}$, we set
$$W_j := \{P \in W \setminus \{0\} : P \text{ is } (1, -1)\text{-homogeneous and } v_{1,-1}(P) = j\} \cup \{0\}.$$

Remark 1.8. It is easy to see that W_j is a subvector space of W. Moreover, by Lemma 1.7, we know that W is a $\mathbb{Z}$-graded algebra with W_j the $(1, -1)$-homogeneous component of degree j, and by [9, Lemma 2.1], we know that $W_0 = K[XY]$, and hence commutative.

Proposition 1.9. *Let* $P, Q \in W \setminus \{0\}$ *and* $(\rho, \sigma) \in \overline{\mathfrak{V}}$. *The following assertions hold:*

1. $w(PQ) = w(P) + w(Q)$ *and* $\overline{w}(PQ) = \overline{w}(P) + \overline{w}(Q)$. *In particular* $PQ \neq 0$.
2. $\ell_{\rho,\sigma}(PQ) = \ell_{\rho,\sigma}(P)\ell_{\rho,\sigma}(Q)$ *for all* $(\rho, \sigma) \in \mathfrak{V}$.
3. $v_{\rho,\sigma}(PQ) = v_{\rho,\sigma}(P) + v_{\rho,\sigma}(Q)$ *for all* $(\rho, \sigma) \in \overline{\mathfrak{V}}$.

(4) $\operatorname{st}_{\rho,\sigma}(PQ)=\operatorname{st}_{\rho,\sigma}(P)+\operatorname{st}_{\rho,\sigma}(Q)$ *for all* $(\rho,\sigma)\in\mathfrak{V}$.
(5) $\operatorname{en}_{\rho,\sigma}(PQ)=\operatorname{en}_{\rho,\sigma}(P)+\operatorname{en}_{\rho,\sigma}(Q)$ *for all* $(\rho,\sigma)\in\mathfrak{V}$.

The same properties hold for $P,Q\in L\setminus\{0\}$.

Proof. For $P,Q\in W\setminus\{0\}$ this follows easily from Lemma 1.7 using that $\rho+\sigma>0$ if $(\rho,\sigma)\in\mathfrak{V}$. The proof for $P,Q\in L\setminus\{0\}$ is easier. $\square$



Illustration of Proposition 1.9 with
$P=X^3Y+X^4Y^3+X^2Y^5+Y^5+XY$
and
$Q=X^3Y^2+X^2Y^3+XY^3+X^2Y^2$.

FIGURE 2

By definition the *cross product* of two vectors $A=(a_1,a_2)$ and $B=(b_1,b_2)$ in $\mathbb{R}^2$ is $A\times B:=\det\left(\begin{smallmatrix}a_1 & a_2\\ b_1 & b_2\end{smallmatrix}\right)$.

Definition 1.10. We say that two vectors A and B in $\mathbb{R}^2$ are *aligned*, if $A\times B=0$.

Definition 1.11. Let $P,Q\in L\setminus\{0\}$. We say that P and Q are *aligned* and write $P\sim Q$, if $w(P)$ and $w(Q)$ are so. Moreover we say that $P,Q\in W\setminus\{0\}$ are aligned if $\Psi(P)\sim\Psi(Q)$. Note that

- By definition $P\sim Q$ if and only if $\ell_{1,-1}(P)\sim\ell_{1,-1}(Q)$.
- $\sim$ is not an equivalence relation (it is so restricted to $\{P: w(P)\neq(0,0)\}$).
- If $P\sim Q$ and $w(P)\neq(0,0)\neq w(Q)$, then $w(P)=\lambda w(Q)$ with $\lambda>0$.

Proposition 1.12. *Let* $P,Q\in W\setminus\{0\}$. *The following assertions hold:*

(1) *If* $P\nsim Q$, *then*
$$[P,Q]\neq 0\quad\textit{and}\quad w\bigl([P,Q]\bigr)=w(P)+w(Q)-(1,1).$$
(2) *If* $\overline{w}(P)\nsim\overline{w}(Q)$, *then*
$$[P,Q]\neq 0\quad\textit{and}\quad \overline{w}\bigl([P,Q]\bigr)=\overline{w}(P)+\overline{w}(Q)-(1,1).$$

Proof. We only prove item (1) since item (2) is similar. Let $w(P)=(r,s)$ and $w(Q)=(u,v)$. Since $\binom{s}{1}\binom{u}{1}-\binom{v}{1}\binom{r}{1}=(r,s)\times(u,v)\neq 0$, using Lemma 1.7 one can check that
$$\ell_t\bigl([P,Q]\bigr)=\left(\binom{s}{1}\binom{u}{1}-\binom{v}{1}\binom{r}{1}\right)\ell_c(P)\ell_c(Q)X^{r+u-1}Y^{s+v-1}.$$
So, $w\bigl([P,Q]\bigr)=w(P)+w(Q)-(1,1)$. $\square$

Remark 1.13. For all $P,Q\in W\setminus\{0\}$ and each $(\rho,\sigma)\in\overline{\mathfrak{V}}$, we have
$$[P,Q]=0\quad\text{or}\quad v_{\rho,\sigma}([P,Q])\leq v_{\rho,\sigma}(P)+v_{\rho,\sigma}(Q)-(\rho+\sigma).$$



$$w\big([P,Q]\big) = w(P) + w(Q) - (1,1)$$

and

$$\overline{w}\big([P,Q]\big) = \overline{w}(P) + \overline{w}(Q) - (1,1).$$

FIGURE 3. Proposition 1.12.

Definition 1.14. Let $(\rho,\sigma) \in \overline{\mathfrak{V}}$ and $P, Q \in W \setminus \{0\}$. We say that P and Q are (ρ,σ)-*proportional* if $[P,Q] = 0$ or $v_{\rho,\sigma}([P,Q]) < v_{\rho,\sigma}(P) + v_{\rho,\sigma}(Q) - (\rho+\sigma)$.

Definition 1.15. For $(\rho,\sigma) \in \overline{\mathfrak{V}}$, we define

$$[-,-]_{\rho,\sigma} \colon \big(W \setminus \{0\}\big) \times \big(W \setminus \{0\}\big) \to L,$$

by

$$[P,Q]_{\rho,\sigma} = \begin{cases} 0 & \text{if } P \text{ and } Q \text{ are } (\rho,\sigma)\text{-proportional},\\ \ell_{\rho,\sigma}([P,Q]) & \text{if } P \text{ and } Q \text{ are not } (\rho,\sigma)\text{-proportional}. \end{cases}$$

Lemma 1.16. *Let $(\rho,\sigma) \in \mathfrak{V}$ and let P and Q be (ρ,σ)-homogeneous elements of $W \setminus \{0\}$.*

1. *If $w(P) \nsim w(Q)$, then $[P,Q] \neq 0$ and $w([P,Q]) = w\big(\ell_{\rho,\sigma}([P,Q])\big)$.*
2. *If $\overline{w}(P) \nsim \overline{w}(Q)$, then $[P,Q] \neq 0$ and $\overline{w}([P,Q]) = \overline{w}\big(\ell_{\rho,\sigma}([P,Q])\big)$.*

Proof. We only prove item (1) since item (2) is similar. Write

$$P = \sum_{i=0}^{\alpha} \lambda_i X^{r-i\sigma} Y^{s+i\rho} \quad \text{and} \quad Q = \sum_{j=0}^{\beta} \mu_j X^{u-j\sigma} Y^{v+j\rho},$$

with $\lambda_0, \lambda_\alpha, \mu_0, \mu_\beta \neq 0$. Since, by Lemma 1.7,

$$X^i Y^j X^{i'} Y^{j'} = \sum_{k=0}^{j} k! \binom{j}{k} \binom{i'}{k} X^{i+i'-k} Y^{j+j'-k},$$

we obtain that

$$[P,Q] = \sum_{i=0}^{\alpha} \sum_{j=0}^{\beta} \sum_{k=0}^{\max\{s+i\rho, v+j\rho\}} \lambda_i \mu_j c_{ijk} X^{r+u-(i+j)\sigma-k} Y^{s+v+(i+j)\rho-k},$$

where

$$c_{ijk} = k! \binom{s+i\rho}{k} \binom{u-j\sigma}{k} - k! \binom{v+j\rho}{k} \binom{r-i\sigma}{k}.$$

Now, since $w(P) \nsim w(Q)$, we have $c_{001} \neq 0$. Consequently, since $\rho+\sigma > 0$,

$$\ell_{\rho,\sigma}([P,Q]) = \sum_{i=0}^{\alpha} \sum_{j=0}^{\beta} \lambda_i \mu_j c_{ij1} x^{r+u-(i+j)\sigma-1} y^{s+v+((i+j)\rho-1}.$$

Using again that $c_{001} \neq 0$, we obtain that

$$w([P,Q]) = (r+u-1, s+v-1) = w\big(\ell_{\rho,\sigma}([P,Q])\big),$$

as desired. $\square$

Proposition 1.17. *Let* $P, Q, R \in W \setminus \{0\}$ *such that* $[P,Q]_{\rho,\sigma} = \ell_{\rho,\sigma}(R)$, *where* $(\rho,\sigma) \in \mathfrak{V}$. *We have:*

1. *If* $\operatorname{st}_{\rho,\sigma}(P) \nsim \operatorname{st}_{\rho,\sigma}(Q)$, *then*
$$\operatorname{st}_{\rho,\sigma}(P) + \operatorname{st}_{\rho,\sigma}(Q) - (1,1) = \operatorname{st}_{\rho,\sigma}(R).$$
2. *If* $\operatorname{en}_{\rho,\sigma}(P) \nsim \operatorname{en}_{\rho,\sigma}(Q)$, *then*
$$\operatorname{en}_{\rho,\sigma}(P) + \operatorname{en}_{\rho,\sigma}(Q) - (1,1) = \operatorname{en}_{\rho,\sigma}(R).$$


Y
$\operatorname{en}_{2,1}(R)$
$\operatorname{en}_{2,1}(P)$
$\ell_{2,1}(R)$
$\operatorname{st}_{2,1}(R)$
$\operatorname{st}_{2,1}(P)$
$\operatorname{st}_{2,1}(Q) = \operatorname{en}_{2,1}(Q)$
X

FIGURE 4. Proposition 1.17.

Proof. We only prove item (1) and leave the proof of item (2), which is similar, to the reader. Let P_1 and Q_1 be (ρ,σ)-homogeneous elements of $W \setminus \{0\}$, such that

$$v_{\rho,\sigma}(P - P_1) < v_{\rho,\sigma}(P_1) \quad \text{and} \quad v_{\rho,\sigma}(Q - Q_1) < v_{\rho,\sigma}(Q_1). \tag{1.1}$$

Since

$$[P,Q] = [P_1,Q_1] + [P_1, Q - Q_1] + [P - P_1, Q],$$

and, by Remark 1.13, we have

$$\begin{aligned} v_{\rho,\sigma}([P_1, Q - Q_1]) &\le v_{\rho,\sigma}(P_1) + v_{\rho,\sigma}(Q - Q_1) - (\rho+\sigma) \\ &< v_{\rho,\sigma}(P_1) + v_{\rho,\sigma}(Q_1) - (\rho+\sigma) \\ &= v_{\rho,\sigma}(P) + v_{\rho,\sigma}(Q) - (\rho+\sigma) \end{aligned}$$

and

$$v_{\rho,\sigma}([P - P_1, Q]) < v_{\rho,\sigma}(P) + v_{\rho,\sigma}(Q) - (\rho+\sigma),$$

it follows from the fact that P and Q are not (ρ,σ)-proportional, that

$$v_{\rho,\sigma}([P,Q] - [P_1,Q_1]) < v_{\rho,\sigma}([P,Q]). \tag{1.2}$$

Note that (1.1) and (1.2) imply

$$\ell_{\rho,\sigma}(P) = \ell_{\rho,\sigma}(P_1), \quad \ell_{\rho,\sigma}(Q) = \ell_{\rho,\sigma}(Q_1) \quad \text{and} \quad \ell_{\rho,\sigma}([P,Q]) = \ell_{\rho,\sigma}([P_1,Q_1]).$$

Consequently, by item (1) of Proposition 1.12 and item (1) of Lemma 1.16,

$$\begin{aligned}\mathrm{st}_{\rho,\sigma}(P)+\mathrm{st}_{\rho,\sigma}(Q)-(1,1) &= \mathrm{st}_{\rho,\sigma}(P_1)+\mathrm{st}_{\rho,\sigma}(Q_1)-(1,1)\\ &= w(P_1)+w(Q_1)-(1,1)\\ &= w([P_1,Q_1])\\ &= w\big(\ell_{\rho,\sigma}([P_1,Q_1])\big)\\ &= w\big(\ell_{\rho,\sigma}([P,Q])\big)\\ &= w\big(\ell_{\rho,\sigma}(R)\big)\\ &= \mathrm{st}_{\rho,\sigma}(R),\end{aligned}$$

as desired. □

Proposition 1.18. *Let $(\rho,\sigma)\in\mathfrak{V}$ and $P,Q\in W\setminus\{0\}$. If*

$$\ell_{\rho,\sigma}(P)=\sum_{i=0}^{\alpha}\lambda_i x^{r-i\sigma}y^{s+i\rho}\quad\text{and}\quad \ell_{\rho,\sigma}(Q)=\sum_{j=0}^{\beta}\mu_j x^{u-j\sigma}y^{v+j\rho},$$

with $\lambda_0,\lambda_\alpha,\mu_0,\mu_\beta\neq 0$, then

$$[P,Q]_{\rho,\sigma}=\sum\lambda_i\mu_j c_{ij}x^{r+u-(i+j)\sigma-1}y^{s+v+(i+j)\rho-1},$$

where $c_{ij}=(u-j\sigma,v+j\rho)\times(r-i\sigma,s+i\rho)$.

Proof. Write

$$P=\sum_{i=0}^{\alpha}\lambda_i X^{r-i\sigma}Y^{s+i\rho}+R_P\quad\text{and}\quad Q=\sum_{j=0}^{\beta}\mu_j X^{u-j\sigma}Y^{v+j\rho}+R_Q.$$

Since $R_P=0$ or $v_{\rho,\sigma}(R_P)<v_{\rho,\sigma}(P)$, and $R_Q=0$ or $v_{\rho,\sigma}(R_Q)<v_{\rho,\sigma}(Q)$, from Remark 1.13 it follows that

$$[P,Q]=\sum_{i=0}^{\alpha}\sum_{j=0}^{\beta}\lambda_i\mu_j\left[X^{r-i\sigma}Y^{s+i\rho},X^{u-j\sigma}Y^{v+j\rho}\right]+R,\tag{1.3}$$

where $R=0$ or $v_{\rho,\sigma}(R)<v_{\rho,\sigma}(P)+v_{\rho,\sigma}(Q)-(\rho+\sigma)$. Now, since $\rho+\sigma>0$ and by Lemma 1.7,

$$X^iY^jX^{i'}Y^{j'}=\sum_{k=0}^{j}k!\binom{j}{k}\binom{i'}{k}X^{i+i'-k}Y^{j+j'-k},$$

we obtain that

$$\left[X^{r-i\sigma}Y^{s+i\rho},X^{u-j\sigma}Y^{v+j\rho}\right]=c_{ij}X^{r+u-(i+j)\sigma-1}Y^{s+v+(i+j)\rho-1}+R_{ij},\tag{1.4}$$

with $R_{ij}=0$ or $v_{\rho,\sigma}(R_{ij})<v_{\rho,\sigma}(P)+v_{\rho,\sigma}(Q)-(\rho+\sigma)$. Combining (1.3) with (1.4), we obtain that

$$[P,Q]=\sum_{i=0}^{\alpha}\sum_{j=0}^{\beta}\lambda_i\mu_j c_{ij}X^{r+u-(i+j)\sigma-1}Y^{s+v+(i+j)\rho-1}+R_{PQ},$$

where $R_{PQ}=0$ or $v_{\rho,\sigma}(R_{PQ})<v_{\rho,\sigma}(P)+v_{\rho,\sigma}(Q)-(\rho+\sigma)$. Since

$$v_{\rho,\sigma}\left(X^{r+u-(i+j)\sigma-1}Y^{s+v+(i+j)\rho-1}\right)=v_{\rho,\sigma}(P)+v_{\rho,\sigma}(Q)-(\rho+\sigma),$$

the result follows immediately. □

Corollary 1.19. *Let $(\rho,\sigma)\in\mathfrak{V}$ and $P,P_1,Q,Q_1\in W\setminus\{0\}$. If $\ell_{\rho,\sigma}(P)=\ell_{\rho,\sigma}(P_1)$ and $\ell_{\rho,\sigma}(Q)=\ell_{\rho,\sigma}(Q_1)$, then $[P,Q]_{\rho,\sigma}=[P_1,Q_1]_{\rho,\sigma}$.*

Proof. Clear from Proposition 1.18. □

Definition 1.20. Given $P \in L \setminus \{0\}$ and $(\rho,\sigma) \in \mathfrak{V}$, we write

$$f_{P,\rho,\sigma} := \sum_{i=0}^{\gamma} a_i x^i \in K[x],$$

if

$$\ell_{\rho,\sigma}(P) = \sum_{i=0}^{\gamma} a_i x^{r-i\sigma} y^{s+i\rho} \quad \text{with } a_0 \neq 0 \text{ and } a_\gamma \neq 0.$$

Now, for $P \in W$ we set $f_{P,\rho,\sigma} := f_{\Psi(P),\rho,\sigma}$. Note that

$$\operatorname{st}_{\rho,\sigma}(P) = (r,s), \quad \operatorname{en}_{\rho,\sigma}(P) = (r-\gamma\sigma, s+\gamma\rho) \tag{1.5}$$

and

$$\ell_{\rho,\sigma}(P) = x^r y^s f_{P,\rho,\sigma}(x^{-\sigma} y^{\rho}). \tag{1.6}$$

Remark 1.21. Let $(\rho,\sigma) \in \mathfrak{V}$. From Proposition 1.9 it follows immediately that

$$f_{PQ,\rho,\sigma} = f_{P,\rho,\sigma} f_{Q,\rho,\sigma} \quad \text{for } P, Q \in W \setminus \{0\}. \tag{1.7}$$

The same result holds for $P, Q \in L \setminus \{0\}$.

Item (2) of the following theorem justify the terminology "(ρ,σ)-proportional" introduced in Definition 1.14.

Theorem 1.22. *Let $(\rho,\sigma) \in \mathfrak{V}$, $P, Q \in W \setminus \{0\}$, $a := v_{\rho,\sigma}(Q)$ and $b := v_{\rho,\sigma}(P)$.*

1. *If $[P,Q]_{\rho,\sigma} \neq 0$, then there exist $h \geq 0$ and $c \in \mathbb{Z}$, such that*
$$x^h f_{[P,Q]} = c f_P f_Q + a x f_P' f_Q - b x f_Q' f_P,$$
where $f_P := f_{P,\rho,\sigma}$, $f_Q := f_{Q,\rho,\sigma}$ and $f_{[P,Q]} := f_{[P,Q],\rho,\sigma}$.
2. *If $[P,Q]_{\rho,\sigma} = 0$ and $a, b > 0$, then there exist $\lambda_P, \lambda_Q \in K^{\times}$, $m, n \in \mathbb{N}$ with* $\gcd(m,n) = 1$ *and a (ρ,σ)-homogeneous polynomial $R \in L$, such that*
$$\frac{m}{n} = \frac{b}{a}, \quad \ell_{\rho,\sigma}(P) = \lambda_P R^m \quad \text{and} \quad \ell_{\rho,\sigma}(Q) = \lambda_Q R^n.$$

Proof. Write

$$\ell_{\rho,\sigma}(P) = \sum_{i=0}^{\alpha} \lambda_i x^{r-i\sigma} y^{s+i\rho} \quad \text{and} \quad \ell_{\rho,\sigma}(Q) = \sum_{j=0}^{\beta} \mu_j x^{u-j\sigma} y^{v+j\rho}, \tag{1.8}$$

with $\lambda_0, \lambda_\alpha, \mu_0, \mu_\beta \neq 0$. By Proposition 1.18,

$$[P,Q]_{\rho,\sigma} = \sum \lambda_i \mu_j c_{ij} x^{r+u-(i+j)\sigma-1} y^{s+v+(i+j)\rho-1},$$

where $c_{ij} := (u-j\sigma, v+j\rho) \times (r-i\sigma, s+i\rho)$. Set

$$F := \sum_{ij} \lambda_i \mu_j c_{ij} x^{i+j}.$$

Note that if $[P,Q]_{\rho,\sigma} = 0$, then $F = 0$, and if $[P,Q]_{\rho,\sigma} \neq 0$, then $F = x^h f_{[P,Q]}$, where h is the multiplicity of x in F. Let

$$a := (u,v) \times (-\sigma,\rho), \quad b := -(-\sigma,\rho) \times (r,s) \quad \text{and} \quad c := (u,v) \times (r,s)$$

Clearly

$$c_{ij} = c + ia - jb, \quad a = v_{\rho,\sigma}(Q) \quad \text{and} \quad b = v_{\rho,\sigma}(P).$$

Since

$$\sum_{i,j} \lambda_i \mu_j x^{i+j} = f_P f_Q, \quad \sum_{i,j} i \lambda_i \mu_j x^{i+j} = x f_P' f_Q \quad \text{and} \quad \sum_{i,j} j \lambda_i \mu_j x^{i+j} = x f_Q' f_P,$$

we obtain

$$F = c f_P f_Q + a x f_P' f_Q - b x f_Q' f_P. \tag{1.9}$$

Item (1) follows immediately from this fact. Assume now that $[P,Q]_{\rho,\sigma}=0$. In this case $F=0$ and, in particular, $c=c_{00}=\frac{F(0)}{\lambda_0\mu_0}=0$. Hence, (1.9) becomes

$$af_P'f_Q-bf_Q'f_P=0.$$

Since $a,b\in\mathbb{N}$, this implies $(f_P^a/f_Q^b)'=0$ and so $f_P^a=\lambda f_Q^b$ for some $\lambda\in K^\times$. Hence, there are $g\in K[x]$ and $\lambda_P,\lambda_Q\in K^\times$, such that

$$f_P=\lambda_P g^m \quad\text{and}\quad f_Q=\lambda_Q g^n, \tag{1.10}$$

where $m:=\frac{b}{\gcd(a,b)}$ and $n:=\frac{a}{\gcd(a,b)}$. Now, note that $\{(s,-r),(\rho,\sigma)\}$ is a basis of $\mathbb{Q}\times\mathbb{Q}$ as a $\mathbb{Q}$-vector space, since

$$(s,-r)\times(\rho,\sigma)=(r,s).(\rho,\sigma)=v_{\rho,\sigma}(P)=a>0,$$

where the dot denotes the usual inner product. Hence, from

$$b(u,v).(s,-r)=b(u,v)\times(r,s)=bc=0=a(r,s).(s,-r)$$

and

$$b(u,v).(\rho,\sigma)=v_{\rho,\sigma}(P)v_{\rho,\sigma}(Q)=v_{\rho,\sigma}(Q)v_{\rho,\sigma}(P)=a(r,s).(\rho,\sigma),$$

it follows that $b(u,v)=a(r,s)$. Consequently $m(u,v)=n(r,s)$, and so there exists $(p,q)\in\mathbb{N}_0\times\mathbb{N}_0$, such that

$$(u,v)=n(p,q) \quad\text{and}\quad (r,s)=m(p,q). \tag{1.11}$$

Write now $g=\sum_{i=0}^{\gamma}\nu_i x^i$ with $\nu_\gamma\neq 0$. Since, by (1.8), (1.10) and (1.11),

$$\gamma m=\alpha,\quad pm=r\quad\text{and}\quad qm=s,$$

we have

$$p-\gamma\sigma=\frac{r-\alpha\sigma}{m}\geq 0\quad\text{and}\quad q+\gamma\rho=\frac{s+\alpha\rho}{m}\geq 0.$$

Consequently

$$R:=\sum_{i=0}^{\gamma}\nu_i x^{p-i\sigma}y^{q+i\rho},$$

is a polynomial which, by (1.10), fulfills

$$\ell_{\rho,\sigma}(P)=x^r y^s f_P(x^{-\sigma}y^{\rho})=\lambda_P\big(x^p y^q g(x^{-\sigma}y^{\rho})\big)^m=\lambda_P R^m$$

and

$$\ell_{\rho,\sigma}(Q)=x^u y^v f_Q(x^{-\sigma}y^{\rho})=\lambda_Q\big(x^p y^q g(x^{-\sigma}y^{\rho})\big)^n=\lambda_Q R^n,$$

as desired. □

Definition 1.23. Let $k\in\mathbb{N}$, $j\in\mathbb{N}_0$, $\varepsilon,b>0$ and $c\in\mathbb{Q}$. We say that a pair (f,g) of polynomials in $K[x]$ satisfies $\mathrm{PE}(k,j,\varepsilon,b,c)$ if there is some $h\in\mathbb{N}_0$, such that

$$x^h f^{k+j}=cf^k g+ax(f^k)'g-bxg'f^k \tag{1.12}$$

is fulfilled, where $a=\frac{j}{k}b+\varepsilon$.

Note that in equation (1.12) each irreducible factor of g that does not divide xf has multiplicity 1.

Proposition 1.24. *If (f,g) satisfy $\mathrm{PE}(k,j,\varepsilon,b,c)$ and $f(0)\neq 0\neq g(0)$, then each irreducible factor u of f, with multiplicity m_u in f, has multiplicity jm_u+1 in g. Consequently $g=f^j\overline{g}$ for some $\overline{g}\in K[x]$ separable and the number of different irreducible factors of f is lower or equal than the degree of $\overline{g}$.*

Proof. We can assume that K is algebraically closed. Take an irreducible monic factor u of f. Since $f(0) \neq 0$, there exists $d \in k^{\times}$ such that $u = x + d$. Write $f = u^s\overline{f}$ and $g = u^r\overline{g}$, with $r, s \in \mathbb{N}$ such that u does not divide $\overline{f}\overline{g}$. Now

$$f' = su^{s-1}u'\overline{f} + u^s\overline{f}' \quad \text{and} \quad g' = ru^{r-1}u'\overline{g} + u^r\overline{g}',$$

and so (1.12) reads

$$x^h\overline{f}^{k+j}u^{s(k+j)} = u^{r+ks-1}(aks - br)xu'\overline{f}^k\overline{g} + u^{r+ks}\overline{f}^{k-1}\big(\overline{f}\overline{g}c + kax\overline{g}\overline{f}' - bx\overline{f}\overline{g}'\big).$$

We claim that $(aks - br) \neq 0$. In fact, on the contrary $s(k+j) \geq r + ks$ and so $sj \geq r$. Since $\varepsilon, b, s, k > 0$, this leads to the contradiction

$$aks - br = \left(\frac{j}{k}b + \varepsilon\right)ks - br = ks\varepsilon + b(js - r) \geq ks\varepsilon > 0.$$

Since u does not divide $xu'\overline{f}^k\overline{g}$, we have $s(k+j) = r + ks - 1$. That is $r = js + 1$, which proves the first assertion. The remainding assertions now follow easily. □

Lemma 1.25. *Let $C, E \in W \setminus \{0\}$, $m \in \mathbb{N}$ and $(\rho,\sigma) \in \mathfrak{V}$. If $[C, E] \neq 0$, then*

$$[C^m, E] \neq 0 \quad \textit{and} \quad \ell_{\rho,\sigma}([C^m, E]) = m\ell_{\rho,\sigma}(C)^{m-1}\ell_{\rho,\sigma}([C, E]).$$

Moreover $[C^m, E]_{\rho,\sigma} \neq 0$ if and only if $[C, E]_{\rho,\sigma} \neq 0$.

Proof. Since

$$[C^m, E] = \sum_{i=0}^{m-1} C^i[C, E]C^{m-i-1}$$

and, by Proposition 1.9,

$$\ell_{\rho,\sigma}(C^i[C, E]C^{m-i-1}) = \ell_{\rho,\sigma}(C)^{m-1}\ell_{\rho,\sigma}([C, E]),$$

we have

$$\ell_{\rho,\sigma}([C^m, E]) = m\ell_{\rho,\sigma}(C)^{m-1}\ell_{\rho,\sigma}([C, E]),$$

In order to prove the last assertion note that, again by Proposition 1.9,

$$v_{\rho,\sigma}([C^m, E]) = v_{\rho,\sigma}(C^m) + v_{\rho,\sigma}([C, E]) - v_{\rho,\sigma}(C),$$

and so

$$v_{\rho,\sigma}([C^m, E]) = v_{\rho,\sigma}(C^m) + v_{\rho,\sigma}(E) - (\rho + \sigma)$$

if and only if

$$v_{\rho,\sigma}([C, E]) = v_{\rho,\sigma}(C) + v_{\rho,\sigma}(E) - (\rho + \sigma),$$

which by Definition 1.15, means that $[C^m, E]_{\rho,\sigma} \neq 0 \Leftrightarrow [C, E]_{\rho,\sigma} \neq 0$. □

Theorem 1.26. *Let $C, D \in W \setminus \{0\}$ with $v_{\rho,\sigma}(C) > 0$, where $(\rho,\sigma) \in \mathfrak{V}$. If*

$$[C^k, D]_{\rho,\sigma} = \ell_{\rho,\sigma}(C^{k+j}) \quad \textit{for some } k \in \mathbb{N} \textit{ and } j \in \mathbb{N}_0, \tag{1.13}$$

then there exists a (ρ,σ)-homogeneous element $E \in W$, such that

$$[C^t, E]_{\rho,\sigma} = t\ell_{\rho,\sigma}(C^t) \quad \textit{for all } t \in \mathbb{N}.$$

Proof. By equation (1.13) and items (1) and (2) of Proposition 1.9, we have

$$(k+j)v_{\rho,\sigma}(C) = kv_{\rho,\sigma}(C) + v_{\rho,\sigma}(D) - (\rho + \sigma) \quad \text{and} \quad \ell_{\rho,\sigma}([C^k, D]) = \ell_{\rho,\sigma}(C^{k+j}),$$

and so,

$$v_{\rho,\sigma}(D) = jv_{\rho,\sigma}(C) + \rho + \sigma \quad \text{and} \quad f_{[C^k,D],\rho,\sigma} = f_{C^{k+j},\rho,\sigma}. \tag{1.14}$$

Hence, by item (1) of Theorem 1.22 and equality (1.7), there exist $h \in \mathbb{N}_0$ and $c \in \mathbb{Z}$, such that

$$x^h f^{k+j} = cf^k g + ax(f^k)'g - bxf^k g',$$


Y
X
$\ell_{\rho,\sigma}(C^3)=[C^2,D]_{\rho,\sigma}$
$\ell_{\rho,\sigma}(C)=[C,E]_{\rho,\sigma}$
$E=\ell_{\rho,\sigma}(E)$

FIGURE 5. Theorem 1.26.

where

$$f:=f_{C,\rho,\sigma},\quad g:=f_{D,\rho,\sigma},\quad a:=v_{\rho,\sigma}(D)\quad\text{and}\quad b:=v_{\rho,\sigma}(C^k)=kv_{\rho,\sigma}(C).$$

Note that, since

$$a=\frac{j}{k}b+\rho+\sigma,$$

the pair (f,g) fulfills $\mathrm{PE}(k,j,\varepsilon,b,c)$, where $\varepsilon:=\rho+\sigma$. By Proposition 1.24 there exists $\overline{g}\in K[x]$ such that $g=f^j\overline{g}$. Set $\alpha:=\deg f$, $\beta:=\deg g$ and write

$$\ell_{\rho,\sigma}(C)=\sum_{i=0}^{\alpha}\lambda_i x^{r-i\sigma}y^{s+i\rho}\quad\text{and}\quad \ell_{\rho,\sigma}(D)=\sum_{i=0}^{\beta}\mu_i x^{u-i\sigma}y^{v+i\rho},$$

with $\lambda_0,\lambda_\alpha,\mu_0,\mu_\beta\neq 0$. By item (2) of Proposition 1.9,

$$\ell_{\rho,\sigma}(C^k)=\ell_{\rho,\sigma}(C)^k=\left(\sum_{i=0}^{\alpha}\lambda_i x^{r-i\sigma}y^{s+i\rho}\right)^k=\sum_{i=0}^{k\alpha}\overline{\lambda}_i x^{kr-i\sigma}y^{ks+i\rho}$$

with each $\overline{\lambda}_i\in K$. Let $\gamma:=\beta-j\alpha$ be the degree of $\overline{g}$ and write $\overline{g}=\sum_{i=0}^{\gamma}\eta_i x^i$. Note that $\eta_0\neq 0$, since $f^j(0)\eta_0=f^j(0)\overline{g}(0)=g(0)\neq 0$. We define

$$E:=k\sum_{i=0}^{\gamma}\eta_i X^{u-jr-i\sigma}Y^{v-js+i\rho}.$$

We claim that $E\in W$. For this it suffices to check that

$$u-jr\geq 0,\quad v-js\geq 0,\quad u-jr-\gamma\sigma\geq 0\quad\text{and}\quad v-js+\gamma\rho\geq 0.$$

In order to prove the two first inequalities, we consider the two cases

$$\mathrm{st}_{\rho,\sigma}(D)\sim\mathrm{st}_{\rho,\sigma}(C)\quad\text{and}\quad\mathrm{st}_{\rho,\sigma}(D)\nsim\mathrm{st}_{\rho,\sigma}(C).$$

Note that $(r,s)\neq(0,0)$, since

$$r\rho+s\sigma=v_{\rho,\sigma}(C)>0,$$

and similarly $(u,v)\neq(0,0)$, since by the first equality in (1.14),

$$u\rho+v\sigma=v_{\rho,\sigma}(D)=jv_{\rho,\sigma}(C)+\rho+\sigma\geq\rho+\sigma>0.$$

Hence, by the last comment in Definition 1.11, if $\mathrm{st}_{\rho,\sigma}(D)\sim\mathrm{st}_{\rho,\sigma}(C)$, then there exists $\lambda>0$ such that

$$(u,v)=\mathrm{st}_{\rho,\sigma}(D)=\lambda\,\mathrm{st}_{\rho,\sigma}(C)=\lambda(r,s).$$

Consequently, again by the first equality in (1.14),

$$jv_{\rho,\sigma}(C)+\rho+\sigma=v_{\rho,\sigma}(D)=v_{\rho,\sigma}(\mathrm{st}_{\rho,\sigma}(D))=\lambda v_{\rho,\sigma}(\mathrm{st}_{\rho,\sigma}(C))=\lambda v_{\rho,\sigma}(C),$$

which implies $\lambda > j$, since $v_{\rho,\sigma}(C) > 0$ and $\rho+\sigma > 0$. But then

$$u - jr = (\lambda - j)r \geq 0 \quad \text{and} \quad v - js = (\lambda - j)s \geq 0.$$

Assume now that $\mathrm{st}_{\rho,\sigma}(D) \nsim \mathrm{st}_{\rho,\sigma}(C)$. Then

$$\mathrm{st}_{\rho,\sigma}(D) \nsim \mathrm{st}_{\rho,\sigma}(C^k),$$

since $\mathrm{st}_{\rho,\sigma}(C^k) = k\,\mathrm{st}_{\rho,\sigma}(C)$ by item (4) of Proposition 1.9. Hence, equality (1.13), Proposition 1.17 and item (4) of Proposition 1.9 yields

$$\mathrm{st}_{\rho,\sigma}(D) + k\,\mathrm{st}_{\rho,\sigma}(C) - (1,1) = (k+j)\,\mathrm{st}_{\rho,\sigma}(C),$$

which implies $(u,v) = j(r,s) + (1,1)$, and so

$$u - jr = 1 > 0 \quad \text{and} \quad v - js = 1 > 0.$$

For the two last inequalities, we consider the two cases

$$\mathrm{en}_{\rho,\sigma}(D) \sim \mathrm{en}_{\rho,\sigma}(C) \quad \text{and} \quad \mathrm{en}_{\rho,\sigma}(D) \nsim \mathrm{en}_{\rho,\sigma}(C).$$

Note that $(r - \alpha\sigma, s + \alpha\rho) \neq (0,0)$, since

$$(r - \alpha\sigma)\rho + (s + \alpha\rho)\sigma = v_{\rho,\sigma}(C) > 0,$$

and similarly $(u - \beta\sigma, v + \beta\rho) \neq (0,0)$, since

$$(u - \beta\sigma)\rho + (v + \beta\rho)\sigma = v_{\rho,\sigma}(D) = jv_{\rho,\sigma}(C) + \rho + \sigma \geq \rho + \sigma > 0.$$

Hence, if $\mathrm{en}_{\rho,\sigma}(D) \sim \mathrm{en}_{\rho,\sigma}(C)$, then arguing as before we obtain $\lambda > 0$ such that

$$(u - \beta\sigma, v + \beta\rho) = \mathrm{en}_{\rho,\sigma}(D) = \lambda\,\mathrm{en}_{\rho,\sigma}(C) = \lambda(r - \alpha\sigma, s + \alpha\rho).$$

Consequently

$$jv_{\rho,\sigma}(C) + \rho + \sigma = v_{\rho,\sigma}(D) = v_{\rho,\sigma}(\mathrm{en}_{\rho,\sigma}(D)) = \lambda v_{\rho,\sigma}(\mathrm{en}_{\rho,\sigma}(C)) = \lambda v_{\rho,\sigma}(C)),$$

which implies $\lambda > j$, since $v_{\rho,\sigma}(C) > 0$ and $\rho + \sigma > 0$. But then

$$u - jr - \gamma\sigma = u - jr - (\beta - j\alpha)\sigma = u - \beta\sigma - j(r - \alpha\sigma) = (\lambda - j)(r - \alpha\sigma) \geq 0$$

and

$$v - js + \gamma\rho = v - js + (\beta - j\alpha)\rho = v + \beta\rho - j(s + \alpha\rho) = (\lambda - j)(s + \alpha\rho) \geq 0.$$

Assume now that $\mathrm{en}_{\rho,\sigma}(D) \nsim \mathrm{en}_{\rho,\sigma}(C)$. Then

$$\mathrm{en}_{\rho,\sigma}(D) \nsim \mathrm{en}_{\rho,\sigma}(C^k),$$

since $\mathrm{en}_{\rho,\sigma}(C^k) = k\,\mathrm{en}_{\rho,\sigma}(C)$ by item (4) of Proposition 1.9. Hence Proposition 1.17 yields

$$\mathrm{en}_{\rho,\sigma}(D) + k\,\mathrm{en}_{\rho,\sigma}(C) - (1,1) = (k+j)\,\mathrm{en}_{\rho,\sigma}(C),$$

which implies $(u - \beta\sigma, v + \beta\rho) = j(r - \alpha\sigma, s + \alpha\rho) + (1,1)$, and so

$$u - jr - \gamma\sigma = 1 > 0 \quad \text{and} \quad v - js + \gamma\rho = 1 > 0.$$

Now, since $\eta_0 \neq 0$ and $\eta_\gamma \neq 0$, we have $\overline{g} = f_{\frac{1}{k}E,\rho,\sigma}$. Consequently, by Proposition 1.9 and the comment in Definition 1.20,

$$\begin{aligned}
\ell_{\rho,\sigma}\left(\frac{1}{k}EC^j\right) &= \ell_{\rho,\sigma}\left(\frac{1}{k}E\right)\ell_{\rho,\sigma}(C)^j \\
&= x^{u-jr}y^{v-js}\overline{g}(x^{-\sigma}y^{\rho})\left(x^r y^s f(x^{-\sigma}y^{\rho})\right)^j \\
&= x^u y^v g(x^{-\sigma}y^{\rho}) \\
&= \ell_{\rho,\sigma}(D).
\end{aligned}$$

Hence, by Corollary 1.19, equality (1.13) and item (2) of Proposition 1.9, we have

$$\left[C^k, \frac{1}{k}EC^j\right]_{\rho,\sigma} = [C^k, D]_{\rho,\sigma} = \ell_{\rho,\sigma}(C^{k+j}) = \ell_{\rho,\sigma}(C)^{k+j}. \tag{1.15}$$

Consequently $\left[C^k, \frac{1}{k}EC^j\right]_{\rho,\sigma} \neq 0$, and so, by items (2) and (3) of Proposition 1.9,

$$\begin{aligned} v_{\rho,\sigma}\left(\left[C^k, \frac{1}{k}EC^j\right]\right) &= v_{\rho,\sigma}(C^k) + v_{\rho,\sigma}\left(\frac{1}{k}EC^j\right) - (\rho+\sigma) \\ &= (k+j)v_{\rho,\sigma}(C) + v_{\rho,\sigma}(E) - (\rho+\sigma) \end{aligned} \tag{1.16}$$

and

$$\begin{aligned} \left[C^k, \frac{1}{k}EC^j\right]_{\rho,\sigma} &= \ell_{\rho,\sigma}\left(\left[C^k, \frac{1}{k}EC^j\right]\right) \\ &= \ell_{\rho,\sigma}\left(\left[C^k, \frac{1}{k}E\right]C^j\right) \\ &= \frac{1}{k}\ell_{\rho,\sigma}([C^k,E])\ell_{\rho,\sigma}(C)^j \\ &= \ell_{\rho,\sigma}([C,E])\ell_{\rho,\sigma}(C)^{k+j-1}, \end{aligned} \tag{1.17}$$

where the last equality follows from Lemma 1.25. Hence, again by Proposition 1.9,

$$v_{\rho,\sigma}\left(\left[C^k, \frac{1}{k}EC^j\right]\right) = (k+j-1)v_{\rho,\sigma}(C) + v_{\rho,\sigma}([C,E]). \tag{1.18}$$

Combining now (1.16) with (1.18), and (1.15) with (1.17), we obtain

$$v_{\rho,\sigma}([C,E]) = v_{\rho,\sigma}(C) + v_{\rho,\sigma}(E) - (\rho+\sigma) \quad \text{and} \quad \ell_{\rho,\sigma}([C,E]) = \ell_{\rho,\sigma}(C).$$

Hence $[C,E]_{\rho,\sigma} = \ell_{\rho,\sigma}(C) \neq 0$, and by Lemma 1.25, we have

$$[C^t,E]_{\rho,\sigma} = \ell_{\rho,\sigma}([C^t,E]) = t\ell_{\rho,\sigma}(C)^{t-1}\ell_{\rho,\sigma}([C,E]) = t\ell_{\rho,\sigma}(C)^t = t\ell_{\rho,\sigma}(C^t),$$

for all $t \in \mathbb{N}$. □

2 The boundary

In this section we establish an order relation on $\overline{\mathfrak{V}}$ and then we describe the boundary of the (convex hull of the) support of an arbitrary element.

We define an order relation on $\mathfrak{V}$ by setting $(\rho_1,\sigma_1) \le (\rho,\sigma)$ if $(\rho_1,\sigma_1) \times (\rho,\sigma) \ge 0$.

To check that $\le$ is indeed an order we first verify that that

$$(\rho_1,\sigma_1) \le (\rho,\sigma) \text{ and } (\rho,\sigma) \le (\rho_1,\sigma_1) \Longrightarrow (\rho_1,\sigma_1) = (\rho,\sigma).$$

In other words that $(\rho_1,\sigma_1) \times (\rho,\sigma) = 0$ if and only if $(\rho_1,\sigma_1) = (\rho,\sigma)$. In fact, if the cross product vanishes, then $(\rho_1,\sigma_1) = \lambda(\rho,\sigma)$ for some $\lambda \in \mathbb{Q} \setminus \{0\}$. If we write $\lambda = m/n$ with m and n relatively prime, then $n(\rho_1,\sigma_1) = m(\rho,\sigma)$ implies that $m|\rho_1$ and $m|\sigma_1$, and so $m = \pm 1$. We also have $n = \pm 1$, since $n|\rho$ and $n|\sigma$. Finally $\rho+\sigma > 0$ and $\rho_1+\sigma_1 > 0$ imply $\lambda = 1$.

We now check the transitivity of $\le$. Assume that

$$(\rho,\sigma) \le (\rho_1,\sigma_1) \quad \text{and} \quad (\rho_1,\sigma_1) \le (\rho_2,\sigma_2).$$

This means

$$\rho\sigma_1 - \sigma\rho_1 \ge 0 \quad \text{and} \quad \rho_1\sigma_2 - \sigma_1\rho_2 \ge 0.$$

Multiplying the first inequality by $\rho_2+\sigma_2 > 0$ and the second one by $\rho+\sigma > 0$, we obtain

$$\rho_2\rho\sigma_1 + \sigma_2\rho\sigma_1 - \rho_2\sigma\rho_1 - \sigma_2\sigma\rho_1 \ge 0 \quad \text{and} \quad \rho\rho_1\sigma_2 + \sigma\rho_1\sigma_2 - \rho\sigma_1\rho_2 - \sigma\sigma_1\rho_2 \ge 0.$$

Summing up the two inequalities we arrive at

$$(\rho_1+\sigma_1)(\rho\sigma_2 - \sigma\rho_2) = \rho_1\rho\sigma_2 + \sigma_1\rho\sigma_2 - \rho_1\sigma\rho_2 - \sigma_1\sigma\rho_2 \ge 0,$$

and so $(\rho,\sigma) \le (\rho_2,\sigma_2)$.

We can extend the order relation to all of $\overline{\mathfrak{V}}$ by setting

$$(1,-1) < (\rho,\sigma) < (-1,1) \quad \text{for all } (\rho,\sigma)\in\mathfrak{V}.$$

Note that if $(\rho,\sigma),(\rho_1\sigma_1)\in\overline{\mathfrak{V}}$ and $\{(\rho,\sigma),(\rho_1\sigma_1)\}\neq\{(1,-1),(-1,1)\}$, then

$$(\rho_1,\sigma_1)<(\rho,\sigma)\iff(\rho_1,\sigma_1)\times(\rho,\sigma)>0.$$



The valuations grow counterclockwise.

FIGURE 6. Order relation in $\overline{\mathfrak{V}}$.

Definition 2.1. Let $P\in W\setminus\{0\}$. We define the set of *valuations associated with* P as

$$\operatorname{Val}(P):=\{(\rho,\sigma)\in\mathfrak{V}: \#\operatorname{Supp}(\ell_{\rho,\sigma}(P))>1\},$$

and we set $\overline{\operatorname{Val}}(P):=\operatorname{Val}(P)\cup\{(1,-1),(-1,1)\}$. We make a similar definition for $P\in L\setminus\{0\}$.



$\overline{\operatorname{Val}}(P)=\{(1,-1),(2,-1),(2,1),(1,3),(1,0),(-1,1)\}.$

FIGURE 7. Definition 2.1.

For each $(r,s)\in\mathbb{Z}\times\mathbb{Z}\setminus\mathbb{Z}(1,1)$ there exists a unique $(\rho,\sigma)\in\mathfrak{V}$ such that $v_{\rho,\sigma}(r,s)=0$. In fact clearly

$$(\rho,\sigma):=\begin{cases}\left(-\frac{s}{d},\frac{r}{d}\right) & \text{if } r-s>0,\\ \left(\frac{s}{d},-\frac{r}{d}\right) & \text{if } r-s<0,\end{cases}$$

where $d:=\gcd(r,s)$, fulfills the required condition, and the uniqueness is evident.

Definition 2.2. For $(r,s)\in\mathbb{Z}\times\mathbb{Z}\setminus\mathbb{Z}(1,1)$, we define $\operatorname{val}(r,s)$ to be the unique $(\rho,\sigma)\in\mathfrak{V}$ such that $v_{\rho,\sigma}(r,s)=0$.

Remark 2.3. Note that if $P\in W\setminus\{0\}$ and $(\rho,\sigma)\in\operatorname{Val}(P)$, then

$$(\rho,\sigma)=\operatorname{val}\bigl(\operatorname{en}_{\rho,\sigma}(P)-\operatorname{st}_{\rho,\sigma}(P)\bigr).$$

Our aim is to prove Proposition 2.8, which geometrically means the following. We start at $w(P)$, draw a ray straight with slope 1, and rotate this ray counter-clockwise until it hits another point in $\operatorname{Supp}(P)$. We break the ray here and continue the counter-clockwise rotation of the ray until it hits a new point in $\operatorname{Supp}(P)$, etcetera. We finish this process when we reach $\overline{w}(P)$. Each straight segment of the border of the support of P are the points where the function $v_{\rho,\sigma}$ takes its maximum, for a uniquely determined $(\rho,\sigma)\in\mathfrak{V}$. Similarly, each corner is the point where the functions $v_{\rho,\sigma}$ take its maximum, for all (ρ,σ) with $(\rho_1,\sigma_1)\ge(\rho,\sigma)\ge(\rho_2,\sigma_2)$, where (ρ_1,σ_1) and (ρ_2,σ_2) are the valuations corresponding to the adjacent sides of the corner. It is also geometrically clear that the endpoint of one side is the starting point of the next.

We fix $P\in W\setminus\{0\}$ and $(\rho,\sigma)\in\mathfrak{V}$. We set $\mathrm{en}:=\mathrm{en}_{\rho,\sigma}(P)$ and $\mathrm{st}:=\mathrm{st}_{\rho,\sigma}(P)$ and we consider the following two sets of valuations

$$\operatorname{Valsup}_{\rho,\sigma}(P):=\{\operatorname{val}((i,j)-\mathrm{en}):(i,j)\in\operatorname{Supp}(P)\text{ and }v_{-1,1}(i,j)>v_{-1,1}(\mathrm{en})\}$$

and

$$\operatorname{Valinf}_{\rho,\sigma}(P):=\{\operatorname{val}((i,j)-\mathrm{st}):(i,j)\in\operatorname{Supp}(P)\text{ and }v_{1,-1}(i,j)>v_{1,-1}(\mathrm{st})\}.$$


Y
Pairs (i,j) involved in the definition of $\operatorname{Valsup}_{\rho,\sigma}(P)$.
en
$\ell_{\rho,\sigma}(P)$
st
Pairs (i,j) involved in the definition of $\operatorname{Valinf}_{\rho,\sigma}(P)$.
X

FIGURE 8. Illustration of the concepts of $\operatorname{Valsup}_{\rho,\sigma}(P)$ and $\operatorname{Valinf}_{\rho,\sigma}(P)$.

Lemma 2.4. *The following assertions hold:*

1. *If* $(\rho_1,\sigma_1)\in\operatorname{Valsup}_{\rho,\sigma}(P)$, *then* $(\rho_1,\sigma_1)>(\rho,\sigma)$.
2. *If* $(\rho_1,\sigma_1)\in\operatorname{Valinf}_{\rho,\sigma}(P)$, *then* $(\rho_1,\sigma_1)<(\rho,\sigma)$.

Proof. We only prove item (1) and leave the other one to the reader. Clearly, if

$$(i,j)\in\operatorname{Supp}(P)\quad\text{and}\quad v_{\rho,\sigma}(i,j)=v_{\rho,\sigma}(P),$$

then $(i,j)\in\operatorname{Supp}\bigl(\ell_{\rho,\sigma}(P)\bigr)$, and so $v_{-1,1}(i,j)\le v_{-1,1}(\mathrm{en})$. Consequently, if

$$(i,j)\in\operatorname{Supp}(P)\quad\text{and}\quad v_{-1,1}(i,j)>v_{-1,1}(\mathrm{en}),$$

then $v_{\rho,\sigma}(i,j)<v_{\rho,\sigma}(P)=v_{\rho,\sigma}(\mathrm{en})$. This means

$$v_{\rho,\sigma}(a,b)<0,\quad\text{where }(a,b):=(i,j)-\mathrm{en}.$$

Note that $v_{-1,1}(i,j)>v_{-1,1}(\mathrm{en})$ now reads

$$b-a=v_{-1,1}(a,b)>0.$$

But then

$$(\rho_1,\sigma_1):=\operatorname{val}\bigl((i,j)-\mathrm{en}\bigr)=\operatorname{val}(a,b)=\lambda(b,-a),$$

for some $\lambda > 0$. Hence

$$\begin{aligned} 0 &> v_{\rho,\sigma}(a,b)\\ &= a\rho + b\sigma\\ &= -\frac{1}{\lambda}(\sigma_1\rho - \rho_1\sigma)\\ &= -\frac{1}{\lambda}(\rho,\sigma)\times(\rho_1,\sigma_1). \end{aligned}$$

This yields $(\rho,\sigma)\times(\rho_1,\sigma_1) > 0$, and so $(\rho_1,\sigma_1) > (\rho,\sigma)$, as desired. ☐

Lemma 2.5. *Let* P, (ρ,σ), st *and* en *be as before. We have:*

1. *If* $(i,j)\in \operatorname{Supp}(P)$, $(\rho_1,\sigma_1) > (\rho,\sigma)$ *and* $v_{-1,1}(i,j) \le v_{-1,1}(\mathrm{en})$, *then*
$$v_{\rho_1,\sigma_1}(i,j) \le v_{\rho_1,\sigma_1}(\mathrm{en}). \tag{2.1}$$
Moreover, if $(\rho_1,\sigma_1)\ne(-1,1)$, *then equality holds if and only if* $(i,j)=\mathrm{en}$.
2. *If* $(i,j)\in \operatorname{Supp}(P)$, $(\rho_1,\sigma_1) < (\rho,\sigma)$ *and* $v_{1,-1}(i,j) \le v_{1,-1}(\mathrm{st})$, *then*
$$v_{\rho_1,\sigma_1}(i,j) \le v_{\rho_1,\sigma_1}(\mathrm{st}).$$
Moreover, if $(\rho_1,\sigma_1)\ne(1,-1)$, *then equality holds if and only if* $(i,j)=\mathrm{st}$.



FIGURE 9. Item (1) of Lemma 2.5.

Proof. We prove item (1) and leave the proof of item (2), which is similar, to the reader. Set $(a,b) := (i,j) - \mathrm{en}$. Then, by the hypothesis,

$$\rho\sigma_1 - \sigma\rho_1 > 0 \quad\text{and}\quad b-a\le 0.$$

Hence

$$b\rho\sigma_1 + \sigma\rho_1 a - a\rho\sigma_1 - b\sigma\rho_1 \le 0, \tag{2.2}$$

and the equality holds if and only if $b=a$. We also know that $v_{\rho,\sigma}(i,j)\le v_{\rho,\sigma}(\mathrm{en})$, which means that $\rho a + \sigma b \le 0$. Since $\rho_1+\sigma_1\ge 0$, we obtain

$$\rho_1\rho a + \sigma_1\sigma b + \rho_1\sigma b + \sigma_1\rho a = (\rho a+\sigma b)(\rho_1+\sigma_1)\le 0. \tag{2.3}$$

Summing up (2.2) and (2.3), we obtain

$$0 \ge \rho\rho_1 a + \sigma\sigma_1 b + \rho\sigma_1 b + \sigma\rho_1 a = (\rho+\sigma)(\rho_1 a + \sigma_1 b),$$

and so $v_{\rho_1,\sigma_1}(a,b)\le 0$, as desired. Moreover, if the equality is true, then (2.2) is also an equality, and so $b=a$. Hence $0 = v_{\rho_1,\sigma_1}(a,a) = (\rho_1+\sigma_1)a$, which implies that $a=0$ or $(\rho_1,\sigma_1)=(-1,1)$. Thus, is $(\rho_1,\sigma_1)\ne(-1,1)$ and equality holds in (2.1), then $(i,j)=\mathrm{en}$. ☐

Definition 2.6. If $\mathrm{Valsup}_{\rho,\sigma}(P) \neq \emptyset$, then we define

$$\mathrm{Succ}_{\rho,\sigma}(P) := \min(\mathrm{Valsup}_{\rho,\sigma}(P))$$

and if $\mathrm{Valinf}_{\rho,\sigma}(P) \neq \emptyset$, then we define

$$\mathrm{Pred}_{\rho,\sigma}(P) := \max(\mathrm{Valinf}_{\rho,\sigma}(P)).$$

Lemma 2.7. *The following assertions hold:*

1. $\mathrm{Succ}_{\rho,\sigma}(P) \in \mathrm{Val}(P)$ *and* $\mathrm{en}_{\rho,\sigma}(P) = \mathrm{st}_{\mathrm{Succ}_{\rho,\sigma}(P)}(P)$.
2. $\mathrm{Pred}_{\rho,\sigma}(P) \in \mathrm{Val}(P)$ *and* $\mathrm{st}_{\rho,\sigma}(P) = \mathrm{en}_{\mathrm{Pred}_{\rho,\sigma}(P)}(P)$.



Y
X
en
st
$\ell_{\rho,\sigma}(P)$
$\mathrm{Succ}_{\rho,\sigma}(P)$
(ρ,σ)
$\mathrm{Pred}_{\rho,\sigma}(P)$

FIGURE 10. Definition 2.6 and Lemma 2.7.

Proof. We only prove (1) since (2) is similar. As above we set $\mathrm{en} := \mathrm{en}_{\rho,\sigma}(P)$. Write $(\rho_1,\sigma_1) := \mathrm{Succ}_{\rho,\sigma}(P)$. By definition, there exists an $(i_0,j_0) \in \mathrm{Supp}(P)$, such that

$$v_{-1,1}(i_0,j_0) > v_{-1,1}(\mathrm{en}) \quad \text{and} \quad (\rho_1,\sigma_1) = \mathrm{val}\big((i_0,j_0) - \mathrm{en}\big).$$

Consequently,

$$(i_0,j_0) \neq \mathrm{en} \quad \text{and} \quad v_{\rho_1,\sigma_1}(\mathrm{en}) = v_{\rho_1,\sigma_1}(i_0,j_0). \tag{2.4}$$

Hence $(\rho_1,\sigma_1) \neq (-1,1)$, since, on the contrary, $v_{\rho_1,\sigma_1}(\mathrm{en}) < v_{\rho_1,\sigma_1}(i_0,j_0)$. We claim that $v_{\rho_1,\sigma_1}(P) = v_{\rho_1,\sigma_1}(\mathrm{en})$, which, by (2.4), proves that $(\rho_1,\sigma_1) \in \mathrm{Val}(P)$. In fact, assume on the contrary that there exists $(i,j) \in \mathrm{Supp}(P)$ with

$$v_{\rho_1,\sigma_1}(i,j) > v_{\rho_1,\sigma_1}(\mathrm{en}) \tag{2.5}$$

By item (1) of Lemmas 2.4 and 2.5, necessarily $v_{-1,1}(i,j) > v_{-1,1}(\mathrm{en})$, and so $(a,b) := (i,j) - \mathrm{en}$ fulfills $b - a > 0$. Hence

$$(\rho_2,\sigma_2) := \mathrm{val}\big((i,j) - \mathrm{en}\big) = \mathrm{val}(a,b) = \lambda(b,-a)$$

with $\lambda > 0$. Now (2.5) leads to

$$\begin{aligned} 0 &< (\rho_1,\sigma_1).(a,b) \\ &= \frac{1}{\lambda}(\rho_2\sigma_1 - \sigma_2\rho_1) \\ &= \frac{1}{\lambda}(\rho_2,\sigma_2) \times (\rho_1,\sigma_1), \end{aligned}$$

which implies that $(\rho_2,\sigma_2) < (\rho_1,\sigma_1)$. But this fact is impossible, since (ρ_1,σ_1) is minimal in $\mathrm{Valsup}_{\rho,\sigma}(P)$ and $(\rho_2,\sigma_2) \in \mathrm{Valsup}_{\rho,\sigma}(P)$. This proves the claim and so $\mathrm{Succ}_{\rho,\sigma}(P) \in \mathrm{Val}(P)$.

Finally we will check that $\mathrm{en} = \mathrm{st}_{\rho_1,\sigma_1}(P)$. For this, it suffices to prove that any $(i,j) \in \mathrm{Supp}(\ell_{\rho_1,\sigma_1}(P))$ fulfills $v_{1,-1}(i,j) \leq v_{1,-1}(\mathrm{en})$ or, equivalently, that

$v_{-1,1}(i,j) \geq v_{-1,1}(\mathrm{en})$. To do this we first note that by item (1) of Lemma 2.4 we have $(\rho_1,\sigma_1) > (\rho,\sigma)$. Since $(i,j) \in \mathrm{Supp}(P)$ and $(\rho_1,\sigma_1) \neq (-1,1)$, from item (1) of Lemma 2.5, it follows that if $v_{-1,1}(i,j) < v_{-1,1}(\mathrm{en})$, then $v_{\rho_1,\sigma_1}(i,j) < v_{\rho_1,\sigma_1}(\mathrm{en})$, which is a contradiction. □

Proposition 2.8. *Let $P \in W \setminus \{0\}$ and let $(\rho_1,\sigma_1) > (\rho_2,\sigma_2)$ consecutive elements in $\overline{\mathrm{Val}}(P)$. The following assertions hold:*

1. *If $(\rho_1,\sigma_1) \in \mathrm{Val}(P)$ and $(\rho_1,\sigma_1) > (\rho,\sigma) \geq (\rho_2,\sigma_2)$, then*
$$(\rho_1,\sigma_1) = \mathrm{Succ}_{\rho,\sigma}(P).$$
2. *If $(\rho_2,\sigma_2) \in \mathrm{Val}(P)$ and $(\rho_1,\sigma_1) \geq (\rho,\sigma) > (\rho_2,\sigma_2)$, then*
$$(\rho_2,\sigma_2) = \mathrm{Pred}_{\rho,\sigma}(P).$$
3. *If $(\rho_1,\sigma_1) > (\rho,\sigma) > (\rho_2,\sigma_2)$, then*
$$\{\mathrm{st}_{\rho_1,\sigma_1}(P)\} = \mathrm{Supp}(\ell_{\rho,\sigma}(P)) = \{\mathrm{en}_{\rho_2,\sigma_2}(P)\}.$$



Y
A
X
(ρ,σ)
(ρ_1,σ_1)
(ρ_2,σ_2)
$A := \mathrm{st}_{\rho_1,\sigma_1}(P) = \mathrm{en}_{\rho_2,\sigma_2}(P)$
$\{A\} = \mathrm{Supp}(\ell_{\rho,\sigma}(P))$

FIGURE 11. Proposition 2.8.

Proof. (1) By item (1) of Lemmas 2.4 and 2.7 it is clear that the existence of $\mathrm{Succ}_{\rho,\sigma}(P)$ implies
$$(\rho,\sigma) < \mathrm{Succ}_{\rho,\sigma}(P) \quad \text{and} \quad \mathrm{Succ}_{\rho,\sigma}(P) \in \mathrm{Val}(P).$$
Hence $(\rho_1,\sigma_1) \leq \mathrm{Succ}_{\rho,\sigma}(P)$. So we must prove that $\mathrm{Succ}_{\rho,\sigma}(P)$ exists and that $(\rho_1,\sigma_1) \geq \mathrm{Succ}_{\rho,\sigma}(P)$. For the existence it suffices to prove that $\mathrm{Valsup}_{\rho,\sigma}(P) \neq \emptyset$. Assume on the contrary that $\mathrm{Valsup}_{\rho,\sigma}(P) = \emptyset$. Then, by definition
$$v_{-1,1}(i,j) \leq v_{-1,1}(\mathrm{en}_{\rho,\sigma})(P) \quad \text{for all } (i,j) \in \mathrm{Supp}(P).$$
Consequently, since $(\rho,\sigma) < (\rho_1,\sigma_1) < (-1,1)$, by item (1) of Lemma 2.5,
$$\mathrm{Supp}(\ell_{\rho_1,\sigma_1}(P)) = \{\mathrm{en}_{\rho,\sigma}(P)\},$$
and so $(\rho_1,\sigma_1) \notin \mathrm{Val}(P)$, which is a contradiction.

Now we prove that $(\rho_1,\sigma_1) \geq \mathrm{Succ}_{\rho,\sigma}(P)$. Since (ρ_1,σ_1) is the minimal element of $\mathrm{Val}(P)$ greater than (ρ,σ), it suffices to prove that there exists no $(\rho_3,\sigma_3) \in \mathrm{Val}(P)$ such that $\mathrm{Succ}_{\rho,\sigma}(P) > (\rho_3,\sigma_3) > (\rho,\sigma)$. In other words that
$$\mathrm{Succ}_{\rho,\sigma}(P) > (\rho_3,\sigma_3) > (\rho,\sigma) \Longrightarrow (\rho_3,\sigma_3) \notin \mathrm{Val}(P).$$
So assume that $\mathrm{Succ}_{\rho,\sigma}(P) > (\rho_3,\sigma_3) > (\rho,\sigma)$ and take $(i,j) \in \mathrm{Supp}(\ell_{\rho_3,\sigma_3}(P))$. We assert that $(i,j) = \mathrm{en}_{\rho,\sigma}(P)$, which shows that $\mathrm{Supp}(\ell_{\rho_3,\sigma_3}(P)) = \{\mathrm{en}_{\rho,\sigma}(P)\}$, and consequently that $(\rho_3,\sigma_3) \notin \mathrm{Val}(P)$. In fact, if $v_{-1,1}(i,j) \leq v_{-1,1}(\mathrm{en}_{\rho,\sigma}(P))$, this follows from item (1) of Lemma 2.5, applied to (ρ_3,σ_3) instead of (ρ_1,σ_1).

Assume now that $v_{-1,1}(i,j) \ge v_{-1,1}(\mathrm{en}_{\rho,\sigma}(P))$. Since, by item (1) of Lemma 2.7, we know that $\mathrm{st}_{\mathrm{Succ}_{\rho,\sigma}(P)}(P) = \mathrm{en}_{\rho,\sigma}(P)$, we have

$$v_{1,-1}(i,j) \le v_{1,-1}(\mathrm{en}_{\rho,\sigma}(P)) = v_{1,-1}\bigl(\mathrm{st}_{\mathrm{Succ}_{\rho,\sigma}(P)}(P)\bigr).$$

Hence, applying item (2) of Lemma 2.5, with $\mathrm{Succ}_{\rho,\sigma}(P)$ instead of (ρ,σ) and (ρ_3,σ_3) instead of (ρ_1,σ_1), and taking into account that $(i,j) \in \mathrm{Supp}(\ell_{\rho_3,\sigma_3}(P))$, we obtain

$$v_{\rho_3,\sigma_3}(i,j) = v_{\rho_3,\sigma_3}\bigl(\mathrm{st}_{\mathrm{Succ}_{\rho,\sigma}(P)}(P)\bigr).$$

Consequently, since $(\rho_3,\sigma_3) \ne (1,-1)$, it follows, again by item (2) of Lemma 2.5, that $(i,j) = \mathrm{st}_{\mathrm{Succ}_{\rho,\sigma}(P)}(P) = \mathrm{en}_{\rho,\sigma}(P)$, which proves the assertion.

(2) It is similar to the proof of item (1).

(3) We first prove that if $(\rho_1,\sigma_1) \in \mathrm{Val}(P)$, then

$$\{\mathrm{st}_{\rho_1,\sigma_1}(P)\} = \mathrm{Supp}(\ell_{\rho,\sigma}(P)).$$

Since $\{\mathrm{en}_{\rho,\sigma}(P)\} = \mathrm{Supp}(\ell_{\rho,\sigma}(P))$, this fact follows from item (1) and item (1) of Lemma 2.7. This conclude the proof of the first equality in the statement when $(\rho_1,\sigma_1) < (-1,1)$. Now, a symmetric argument shows that if $(\rho_2,\sigma_2) > (1,-1)$, then

$$\{\mathrm{en}_{\rho_2,\sigma_2}(P)\} = \mathrm{Supp}(\ell_{\rho,\sigma}(P)).$$

Assume now that $(\rho_1,\sigma_1) = (-1,1)$ and $(\rho_2,\sigma_2) \ne (1,-1)$. Then, by item (1) of Lemmas 2.4 and 2.7, we have $\mathrm{Valsup}(\rho_2,\sigma_2) = \emptyset$. Hence

$$v_{-1,1}(i,j) \le v_{-1,1}(\mathrm{en}_{\rho_2,\sigma_2}(P)),$$

for all $(i,j) \in \mathrm{Supp}(P)$. Consequently, $\mathrm{en}_{\rho_2,\sigma_2}(P) \in \mathrm{Supp}\bigl(\ell_{-1,1}(P)\bigr)$, and so

$$\mathrm{st}_{-1,1}(P) = \mathrm{en}_{\rho_2,\sigma_2}(P) + (a,a),$$

for some $a \ge 0$. But necessarily $a = 0$, since $a > 0$ leads to the contradiction

$$v_{\rho_2,\sigma_2}(\mathrm{st}_{-1,1}(P)) = v_{\rho_2,\sigma_2}(\mathrm{en}_{\rho_2,\sigma_2}(P) + (a,a)) = v_{\rho_2,\sigma_2}(P) + a(\rho_2+\sigma_2).$$

Thus

$$\{\mathrm{st}_{\rho_1,\sigma_1}(P)\} = \{\mathrm{en}_{\rho_2,\sigma_2}(P)\} = \mathrm{Supp}(\ell_{\rho,\sigma}(P)).$$

Similarly, if $(\rho_1,\sigma_1) \ne (-1,1)$ and $(\rho_2,\sigma_2) = (1,-1)$, then

$$\{\mathrm{st}_{\rho_1,\sigma_1}(P)\} = \{\mathrm{en}_{\rho_2,\sigma_2}(P)\} = \mathrm{Supp}(\ell_{\rho,\sigma}(P)).$$

Finally we assume that $(\rho_1,\sigma_1) = (-1,1)$ and $(\rho_2,\sigma_2) = (1,-1)$. Since $\mathrm{Val}(P) = \emptyset$, it follows from Lemma 2.7 that

$$\mathrm{Valsup}_{\rho,\sigma}(P) = \emptyset = \mathrm{Valinf}_{\rho,\sigma}(P).$$

Hence

$$v_{-1,1}(P) = v_{-1,1}(\mathrm{en}_{\rho,\sigma}(P)) \quad \text{and} \quad v_{1,-1}(P) = v_{1,-1}(\mathrm{st}_{\rho,\sigma}(P)). \tag{2.6}$$

But, since $\mathrm{en}_{\rho,\sigma}(P) = \mathrm{st}_{\rho,\sigma}(P)$, it follows easily from (2.6) that $P = \ell_{-1,1}(P)$, and so,

$$\{\mathrm{en}_{1,-1}(P)\} = \{w(P)\} = \{\overline{w}(P)\} = \{\mathrm{st}_{-1,1}(P)\} = \mathrm{Supp}\bigl(\ell_{\rho,\sigma}(P)\bigr),$$

as desired. □

3 Irreducible pairs

The aim of this section is to introduce the concept of irreducible pairs and to establish some of its basic properties.

Definition 3.1. We say that an endomorphism ψ of W is *irreducible* if

$$v_{1,1}\big(\psi(X)\big) \ge 2, \quad v_{1,1}\big(\psi(Y)\big) \ge 2$$

and there are no $\phi_1, \phi_2 \in \operatorname{Aut}(W)$ such that

$$v_{1,1}\big(\phi_1 \circ \psi \circ \phi_2(X)\big) + v_{1,1}\big(\phi_1 \circ \psi \circ \phi_2(Y)\big) < v_{1,1}\big(\psi(X)\big) + v_{1,1}\big(\psi(Y)\big).$$

A pair (P, Q) of elements of W is said to be *irreducible* if there is an irreducible endomorphism ψ of W such that $\psi(X) = P$ and $\psi(Y) = Q$. Note that this implies that $[Q, P] = 1$.

Remark 3.2. Clearly if (Q, P) is irreducible, then $(P + \lambda_P, Q + \lambda_Q)$ is also for all $\lambda_P, \lambda_Q \in K$. In fact, it is obvious that

$$v_{1,1}(P + \lambda_P) = v_{1,1}(P) \quad \text{and} \quad v_{1,1}(Q + \lambda_Q) = v_{1,1}(Q)$$

since $v_{1,1}(P) > 0$ and $v_{1,1}(Q) > 0$. In particular

$$v_{1,1}(P + \lambda_P) \ge 2 \quad \text{and} \quad v_{1,1}(Q + \lambda_Q) \ge 2.$$

Now, let ψ and ψ' be the endomorphisms of W defined by $\psi(X) = P$, $\psi(Y) = Q$ and $\psi'(X) = P + \lambda_P$, $\psi'(Y) = \lambda_Q$, respectively. There are no $\phi_1, \phi_2 \in \operatorname{Aut}(W)$, such that

$$v_{1,1}\big(\phi_1 \circ \psi' \circ \phi_2(X)\big) + v_{1,1}\big(\phi_1 \circ \psi' \circ \phi_2(Y)\big) < v_{1,1}\big(\psi'(X)\big) + v_{1,1}\big(\psi'(Y)\big),$$

since this implies that

$$v_{1,1}\big(\phi_1 \circ \psi \circ \phi \circ \phi_2(X)\big) + v_{1,1}\big(\phi_1 \circ \psi \circ \phi \circ \phi_2(Y)\big) < v_{1,1}\big(\psi(X)\big) + v_{1,1}\big(\psi(Y)\big),$$

where $\phi \in \operatorname{Aut}(W)$ is defined by $\phi(X) = X + \lambda_P$ and $\phi(Y) = Y + \lambda_P$, which contradicts the fact that (P, Q) is irreducible.

Theorem 3.3. *If there is no irreducible endomorphism, then every endomorphism of W is an automorphism.*

Proof. Assume we have an endomorphism ψ of W, which is not an automorphism. Let $P = \psi(X)$ and $Q = \psi(Y)$. Suppose we have chosen ψ with $v_{1,1}(P) + v_{1,1}(Q)$ minimal. Since, by hypothesis, ψ is reducible, then necessarily $v_{1,1}(P) = 1$ or $v_{1,1}(Q) = 1$. We can assume that $v_{1,1}(P) = 1$. Hence $P = \lambda_0 + \lambda_1 X + \lambda_2 Y$ with $\lambda_1 \ne 0$ or $\lambda_2 \ne 0$. Let ϕ be the automorphism of W defined by

$$\phi(X) = P \quad \text{and} \quad \phi(Y) = \begin{cases} \lambda_1^{-1} Y & \text{if } \lambda_1 \ne 0, \\ -\lambda_2^{-1} X & \text{if } \lambda_1 = 0. \end{cases}$$

Clearly $\phi^{-1} \circ \psi(X) = X$, and so

$$[\phi^{-1} \circ \psi(Y), X] = [\phi^{-1} \circ \psi(Y), \phi^{-1} \circ \psi(X)] = 1.$$

But this implies that $\phi^{-1} \circ \psi(Y) - Y$ belongs to the centralizer of X, which is $K[X]$. Hence, there exists $p \in K[X]$ such that $\phi^{-1} \circ \psi(Y) = Y + p(X)$. From these follows easily that $\phi^{-1} \circ \psi$ is an automorphism. Consequently ψ is also an automorphism, which is a contradiction. ☐

Next we begin the study of the irreducible pairs. We will need the following two lemmas.

Lemma 3.4. *Let* $(\rho,\sigma)\in\mathfrak{V}$ *and* $P,Q\in W\setminus\{0\}$. *If*

$$\operatorname{Supp}(\ell_{\rho,\sigma}(P))=\{(i,j)\}\quad\textit{and}\quad\operatorname{Supp}(\ell_{\rho,\sigma}(Q))=\{(u,v)\},$$

with $(i,j)\nsim(u,v)$, *then*

$$\operatorname{Supp}(\ell_{\rho,\sigma}([Q,P]))=\{(i+j-1,u+v-1)\}.$$

Proof. By Proposition 1.18. ☐

Lemma 3.5. *If* $P,Q\in W$ *satisfy* $[Q,P]=1$ *and* $w(P)=(0,1)$, *then* $P=\mu Y$ *with* $\mu\in K^{\times}$.

Proof. Note that $P\nsim Q$, since $P\sim Q$ leads to $w(Q)=(0,k)$ with $k\in\mathbb{N}_0$. But then $v_{1,-1}(Q)\le 0$, which by Remark 1.13, implies $v_{1,-1}([Q,P])<0$, a contradiction, since $[Q,P]=1$. Consequently, by Proposition 1.12, we know that $w(Q)=(1,0)$. Now, we will assume that $\operatorname{Supp}(P)\neq\{(0,1)\}$ and we will arrive at a contradiction. Since $(0,1)=w(P)$ we have $\operatorname{Supp}(\ell_{1,-1}(P))=\{(0,1)\}$. Hence $P\neq\ell_{1,-1}(P)$ and so $\operatorname{Val}(P)\neq\emptyset$. Take $(\rho,\sigma):=\min\operatorname{Val}(P)$ and set $A:=\operatorname{en}_{\rho,\sigma}(P)$. By Proposition 2.8 we know that

$$(0,1)=w(P)=\operatorname{en}_{1,-1}(P)=\operatorname{st}_{\rho,\sigma}(P),$$

and so

$$A\neq(0,1)\quad\text{and}\quad v_{\rho,\sigma}(A)=v_{\rho,\sigma}(0,1)=\sigma.$$

We claim that $(\rho,\sigma)\le(1,0)$, or, equivalently, that $\sigma\le 0$, and hence $\rho>0$. In fact, each $(i,j)\in\operatorname{Supp}(P)\setminus\{(0,1)\}$, satisfies $j-i>1$, since $w(P)=(0,1)$. In particular

$$A=(i,i+k)\quad\text{with } i\ge 0 \text{ and } k>1. \tag{3.1}$$

Then

$$\rho i+\sigma(i+k)=v_{\rho,\sigma}(A)=v_{\rho,\sigma}(0,1)=\sigma.$$

Hence, $\sigma(1-k)=i(\rho+\sigma)\ge 0$, which implies $\sigma\le 0$, since $k>1$, proving the claim.



FIGURE 12. Shape of the hypothetical P with $\operatorname{Supp}(P)\neq\{(0,1)\}$.

Now set $B:=\operatorname{en}_{\rho,\sigma}(Q)$. Note that $A\neq(0,0)$, since

$$v_{1,-1}(P)=v_{1,-1}(w(P))=v_{1,-1}(0,1)=-1,$$

and $B\neq(0,0)$, since

$$v_{\rho,\sigma}(B)=v_{\rho,\sigma}(Q)\ge v_{\rho,\sigma}(1,0)=\rho>0.$$

We claim that $A\nsim B$. On the contrary, since $A\neq(0,0)$ and $B\neq(0,0)$, there is $\lambda>0$ such that $\lambda B=A$. But this implies

$$v_{\rho,\sigma}(A)=\lambda v_{\rho,\sigma}(B)>0,$$

which contradicts that $v_{\rho,\sigma}(A)=\sigma\le 0$. Let $(\rho_1,\sigma_1)\in\mathfrak{V}$, such that

- $(\rho_1,\sigma_1) > (\rho,\sigma)$,
- $(\rho_1,\sigma_1) < \min\{((\rho',\sigma') \in \overline{\mathrm{Val}}(P)$ such that $(\rho',\sigma') > (\rho,\sigma)\}$,
- $(\rho_1,\sigma_1) < \min\{((\rho',\sigma') \in \overline{\mathrm{Val}}(Q)$ such that $(\rho',\sigma') > (\rho,\sigma)\}$.

By Proposition 2.8,

$$\mathrm{Supp}(\ell_{\rho_1,\sigma_1}(P)) = \{A\} \quad \text{and} \quad \mathrm{Supp}(\ell_{\rho_1,\sigma_1}(Q)) = \{B\}.$$

Since $A \nsim B$, by Lemma 3.4 we have $A+B = (1,1)$. But this is impossible, because of (3.1). $\square$

Proposition 3.6. *If ψ is irreducible, then $v_{\rho,\sigma}\big(\psi(X)\big) > 0$ and $v_{\rho,\sigma}\big(\psi(Y)\big) > 0$ for all $(\rho,\sigma) \in \overline{\mathfrak{V}}$.*

Proof. Let $(\rho,\sigma) \in \overline{\mathfrak{V}}$, $P = \psi(X)$ and $Q = \psi(Y)$. We will prove that $v_{\rho,\sigma}(P) > 0$ and leave the proof that $v_{\rho,\sigma}(Q) > 0$, which is similar, to the reader. We divide the proof in three cases.

First case. $(1,-1) \le (\rho,\sigma) \le (1,0)$. By definition this means that $\rho \ge -\sigma \ge 0$. Since $\rho = \sigma = 0$ is impossible, necessarily $\rho > 0$. Suppose that $v_{\rho,\sigma}(P) \le 0$. We claim that $v_{1,-1}(P) \le 0$ and $v_{1,-1}(Q) > 0$. The first inequality follows from the fact that $v_{1,-1}(X^iY^j) = i-j \le 0$ for all $(i,j) \in \mathrm{Supp}(P)$, since

$$\begin{aligned}\rho(i-j) + (\rho+\sigma)j = \rho i + \sigma j = v_{\rho,\sigma}(X^iY^j) \le 0 &\Rightarrow \rho(i-j) \le 0\\ &\Rightarrow v_{1,-1}(X^iY^j) = i-j \le 0.\end{aligned}$$

It remains to prove that $v_{1,-1}(Q) > 0$. In fact, otherwise $v_{1,-1}([Q,P]) < 0$ (which contradicts $[Q,P] = 1$), since $P_i = Q_i = 0$ for $i > 0$ and $[Q_0,P_0] = 0$ by Remark 1.8, where P_i and Q_i denote the i-homogeneous components of P and Q, respectively. We assert that

$$P = \lambda_0 + \overline{P} \quad \text{with } \lambda_0 \in K \text{ and } \overline{P} \not\sim Q. \tag{3.2}$$

If $P \nsim Q$, then we can take $\lambda_0 = 0$. Hence, we can assume that $P \sim Q$, which combined with $v_{1,-1}(P) \le 0$ and $v_{1,-1}(Q) > 0$ gives $w(P) = (0,0)$. Consequently $P = \lambda_0 + \overline{P}$ with $\lambda_0 \in K^\times$ and $v_{1,-1}(\overline{P}) < 0$. This implies that $\overline{P} \nsim Q$ and hence $w(\overline{P}) = (0,1)$ (since $v_{1,-1}(\overline{P}) < 0$ and $w(\overline{P}) + w(Q) = (1,1)$ by item (1) of Proposition 1.12). Now, by Lemma 3.5 we have $v_{1,1}(\overline{P}) = 1$ and so $(\overline{P},Q)$ is not irreducible. On the other hand we know by Remark 3.2 that $(\overline{P},Q)$ is irreducible, which leads to the desired contradiction.

Second case. $(1,0) < (\rho,\sigma) < (0,1)$. This means that $\rho,\sigma > 0$. It is evident that $v_{\rho,\sigma}(A) \ge 0$ for all $A \in W \setminus \{0\}$, and that the equality holds if and only if $A \in K^\times$. Consequently, $v_{\rho,\sigma}(P) > 0$, since $[Q,P] = 1$ implies $P \notin K^\times$.

Third case. $(0,1) \le (\rho,\sigma) \le (-1,1)$. This means that $\sigma \ge -\rho \ge 0$ and the proof is similar to the one of the first case. $\square$

Proposition 3.7. *Let (P,Q) be an irreducible pair and let $(\rho,\sigma) \in \overline{\mathfrak{V}}$. Then*

$$[P,Q]_{\rho,\sigma} = 0.$$

Proof. We will prove that for all $(\rho,\sigma) \in \overline{\mathfrak{V}}$,

$$0 < v_{\rho,\sigma}(P) + v_{\rho,\sigma}(Q) - (\rho+\sigma). \tag{3.3}$$

This will prove the proposition, since $v_{\rho,\sigma}([P,Q]) = 0$. If $(\rho,\sigma) \in \{(1,-1),(-1,1)\}$, then Proposition 3.6 guarantees (3.3). Hence we can assume $\rho+\sigma > 0$ and will prove (3.3) in the four cases

$$(\rho,\sigma) \le (1,0), \quad (1,0) < (\rho,\sigma) \le (1,1), \quad (1,1) < (\rho,\sigma) < (0,1), \quad (0,1) \le (\rho,\sigma).$$

If $(\rho,\sigma)\le(1,0)$, then $\sigma\le 0$ and so $\rho>0$. Since, by Proposition 3.6 we have $v_{1,-1}(P)>0$, there exists $(i,j)\in\operatorname{Supp}(P)$ such that $i-j>0$. But then

$$v_{\rho,\sigma}(P)\ge \rho i+\sigma j=(\rho+\sigma)j+(i-j)\rho\ge(i-j)\rho\ge\rho\ge\rho+\sigma.$$

Similarly $v_{\rho,\sigma}(Q)\ge\rho+\sigma$, and so (3.3) follows, since $\rho+\sigma>0$.

If $(1,0)<(\rho,\sigma)\le(1,1)$, then $\rho\ge\sigma>0$. Since $v_{1,1}(P)\ge 2$, there is a pair $(r,s)\in\operatorname{Supp}(P)$, such that $r+s\ge 2$, and so

$$v_{\rho,\sigma}(P)\ge r\rho+s\sigma=(r+s)\sigma+r(\rho-\sigma)\ge(r+s)\sigma\ge 2\sigma. \tag{3.4}$$

Moreover, since $v_{1,0}(Q)>0$ by Proposition 3.6, there exists $(i,j)\in\operatorname{Supp}(Q)$ with $i>0$, and so

$$v_{\rho,\sigma}(Q)\ge i\rho+j\sigma\ge\rho. \tag{3.5}$$

Combining (3.4) and (3.5), we obtain

$$v_{\rho,\sigma}(P)+v_{\rho,\sigma}(Q)\ge\rho+2\sigma>\rho+\sigma.$$

The third case is similar to the second case and the fourth case similar to the first case. $\square$

Proposition 3.8. *If ψ is irreducible, then*

$$\gcd\bigl(v_{1,1}(P),v_{1,1}(Q)\bigr)\notin\{v_{1,1}(P),v_{1,1}(Q)\}.$$

where $P:=\psi(X)$ and $Q:=\psi(Y)$.

Proof. If this false, then

$$v_{1,1}(P)\mid v_{1,1}(Q)\quad\text{or}\quad v_{1,1}(Q)\mid v_{1,1}(P).$$

Assume that $v_{1,1}(P)\mid v_{1,1}(Q)$. By Propositions 3.6 and 3.7, the hypothesis of item (2) of Theorem 1.22 are fulfilled. Hence there exist $\lambda_P,\lambda_Q\in K^\times$, $m,n\in\mathbb{N}$ with $\gcd(m,n)=1$ and a $(1,1)$-homogeneous polynomial $R\in L$, such that

$$\frac{m}{n}=\frac{\operatorname{val}_{1,1}(P)}{\operatorname{val}_{1,1}(Q)},\quad \ell_{1,1}(P)=\lambda_P R^m\quad\text{and}\quad \ell_{1,1}(Q)=\lambda_Q R^n.$$

Consequently $m=1$ and

$$\ell_{1,1}(Q)=\lambda_Q\Bigl(\frac{\ell_{1,1}(P)}{\lambda_P}\Bigr)^n=\lambda\ell_{1,1}(P)^n=\ell_{1,1}(\lambda P^n), \tag{3.6}$$

where $\lambda=\lambda_Q/\lambda_P^n$ and the last equality follows from Proposition 1.9. Let ϕ be the automorphism of W defined by $\phi(X)=X$ and $\phi(Y)=Y-\lambda X^n$. We have

$$v_{1,1}\bigl(\psi\circ\phi(X)\bigr)=v_{1,1}\bigl(\psi(X)\bigr)$$

and

$$v_{1,1}\bigl(\psi\circ\phi(Y)\bigr)=v_{1,1}\bigl(Q-\lambda P^n\bigr)<v_{1,1}(Q)=v_{1,1}\bigl(\psi(Y)\bigr),$$

where the inequality holds by (3.6), which contradicts the fact that the endomorphism ψ is irreducible. $\square$

Remark 3.9. If (P,Q) is an irreducible pair, then $\operatorname{Val}(P)=\operatorname{Val}(Q)$. In fact, if $(\rho,\sigma)\in\mathfrak{V}$, then by Proposition 3.6, Proposition 3.7 and item (2) of Theorem 1.22 there exist $\lambda_P,\lambda_Q\in K^\times$, $m,n\in\mathbb{N}$ and a (ρ,σ)-homogeneous polynomial $R\in L$, such that $\ell_{\rho,\sigma}(P)=\lambda_P R^m$ and $\ell_{\rho,\sigma}(Q)=\lambda_Q R^n$. The assertion follows from the evident fact that

$$\begin{aligned}(\rho,\sigma)\in\operatorname{Val}(P)&\Leftrightarrow \ell_{\rho,\sigma}(P)\text{ is not a monomial}\\&\Leftrightarrow R\text{ is not a monomial}\\&\Leftrightarrow \ell_{\rho,\sigma}(Q)\text{ is not a monomial}\\&\Leftrightarrow(\rho,\sigma)\in\operatorname{Val}(Q).\end{aligned}$$

Consequently, by item (1) of Lemma 2.7 and item (1) of Proposition 2.8, for all $(\rho,\sigma)\in\mathfrak{V}$, we have

$$\operatorname{Succ}_{\rho,\sigma}(P)\text{ exists}\Longrightarrow\operatorname{Succ}_{\rho,\sigma}(Q)\text{ exists and }\operatorname{Succ}_{\rho,\sigma}(Q)=\operatorname{Succ}_{\rho,\sigma}(P);$$

and similarly for $\operatorname{Pred}_{\rho,\sigma}$.

We resume the properties of irreducible pairs (P,Q).

Proposition 3.10. *Let (P,Q) be an irreducible pair and $(\rho,\sigma)\in\mathfrak{V}$. Then we can find a (ρ,σ)-homogeneous element $C\in W$, $\lambda_P,\lambda_Q\in K^{\times}$ and $n,m\in\mathbb{N}$ such that*

1. *$v_{\rho,\sigma}(C)>0$ and $\ell_{\rho,\sigma}(C)\neq\zeta\ell_{\rho,\sigma}(D^h)$ for all $D\in W$, $\zeta\in K^{\times}$ and $h>1$,*
2. *$\ell_{\rho,\sigma}(P)=\lambda_P\ell_{\rho,\sigma}(C^m)$,*
3. *$\ell_{\rho,\sigma}(Q)=\lambda_Q\ell_{\rho,\sigma}(C^n)$,*
4. *$\gcd(n,m)\notin\{n,m\}$.*

Moreover

$$\frac{m}{n}=\frac{v_{\rho,\sigma}(P)}{v_{\rho,\sigma}(Q)}=\frac{v_{1,1}(P)}{v_{1,1}(Q)}. \tag{3.7}$$



FIGURE 13. Proposition 3.10.

Proof. By Proposition 3.6, we know that

$$v_{\rho',\sigma'}(P)>0\quad\text{and}\quad v_{\rho',\sigma'}(Q)>0$$

for all $(\rho',\sigma')\in\overline{\mathfrak{V}}$. We will use again and again this fact freely throughout this proof. Arguing as in the proof of Proposition 3.8 we find $\bar\lambda_P,\bar\lambda_Q\in K^{\times}$, $\bar m,\bar n\in\mathbb{N}$ and a (ρ,σ)-homogeneous polynomial $\bar R\in L$, such that

$$\ell_{\rho,\sigma}(P)=\bar\lambda_P\bar R^{\bar m}\quad\text{and}\quad\ell_{\rho,\sigma}(Q)=\bar\lambda_Q\bar R^{\bar n}.$$

Write

$$\bar R=\lambda R^k\quad\text{with }\lambda\in K^{\times}\text{ and }k\text{ maximal.}$$

Let $C:=\Psi^{-1}(R)$, where Ψ is the map introduced at the beginning of Section 1. Since $\ell_{\rho,\sigma}(C)=R$, it follows immediately from Proposition 1.9 that C fulfills the second part of item (1). Now, since

$$\ell_{\rho,\sigma}(P)=\bar\lambda_P\bar R^{\bar m}=\bar\lambda_P\lambda^kR^{k\bar m}\quad\text{and}\quad\ell_{\rho,\sigma}(Q)=\bar\lambda_Q\bar R^{\bar n}=\bar\lambda_Q\lambda^kR^{k\bar n},$$

the same proposition shows that C fulfills items (2) and (3) with

$$\lambda_P=\bar\lambda_P\lambda^k,\quad\lambda_Q=\bar\lambda_Q\lambda^k,\quad m=k\bar m\quad\text{and}\quad n=k\bar n.$$

Hence $v_{\rho,\sigma}(C)=\frac{1}{m}v_{\rho,\sigma}(P)>0$, and so the first part of item (1) is also fulfilled. Since

$$v_{\rho,\sigma}(P)=mv_{\rho,\sigma}(R)\quad\text{and}\quad v_{\rho,\sigma}(Q)=nv_{\rho,\sigma}(R), \tag{3.8}$$

we have $m/n=v_{\rho,\sigma}(P)/v_{\rho,\sigma}(Q)$. It remains to prove that item (4) also holds. By (3.8) and Proposition 3.8 it suffices to check equality (3.7). In order to do this it is sufficient to show that if

$$(\rho_1,\sigma_1)<(\rho',\sigma')<(\rho_2,\sigma_2),$$

where (ρ_1,σ_1) and (ρ_2,σ_2) are consecutive elements in $\mathrm{Val}(P)=\mathrm{Val}(Q)$, then

$$\frac{v_{\rho',\sigma'}(P)}{v_{\rho',\sigma'}(Q)}=\frac{v_{\rho_1,\sigma_1}(P)}{v_{\rho_1,\sigma_1}(Q)}=\frac{v_{\rho_2,\sigma_2}(P)}{v_{\rho_2,\sigma_2}(Q)}. \tag{3.9}$$

Arguing again as in the proof of Proposition 3.8 we find $\lambda'_P,\lambda'_Q\in K^{\times}$, $m',n'\in\mathbb{N}$ and a (ρ',σ')-homogeneous polynomial $R'\in L$, such that

$$\ell_{\rho',\sigma'}(P)=\lambda'_P R'^{m'}\quad\text{and}\quad \ell_{\rho',\sigma'}(Q)=\lambda'_Q R'^{n'}.$$

Note that R' is a monomial since $(\rho',\sigma')\notin\mathrm{Val}(P)$. Write $\mathrm{Supp}(R')=\{(a,b)\}$. By Proposition 2.8,

$$\{\mathrm{en}_{\rho_1,\sigma_1}(P)\}=\{\mathrm{st}_{\rho_2,\sigma_2}(P)\}=\mathrm{Supp}(\ell_{\rho',\sigma'}(P))=\{m'(a,b)\}$$

and

$$\{\mathrm{en}_{\rho_1,\sigma_1}(Q)\}=\{\mathrm{st}_{\rho_2,\sigma_2}(Q)\}=\mathrm{Supp}(\ell_{\rho',\sigma'}(Q))=\{n'(a,b)\}.$$

Hence

$$\begin{aligned}
v_{\rho',\sigma'}(P)&=m'(\rho'a+\sigma'b), & v_{\rho',\sigma'}(Q)&=n'(\rho'a+\sigma'b),\\
v_{\rho_1,\sigma_1}(P)&=m'(\rho_1a+\sigma_1b), & v_{\rho_1,\sigma_1}(Q)&=n'(\rho_1a+\sigma_1b),\\
v_{\rho_2,\sigma_2}(P)&=m'(\rho_2a+\sigma_2b), & v_{\rho_2,\sigma_2}(Q)&=n'(\rho_2a+\sigma_2b),
\end{aligned}$$

from which (3.9) follows immediately. □

Remark 3.11. Note that from items (1) and (2) of Proposition 3.10, Remark 3.6 and items (4) and (5) of Proposition 1.9, it follows that, for any irreducible pair (P,Q) and any $(\rho,\sigma)\in\mathfrak{V}$,

$$\begin{aligned}
&\mathrm{st}_{\rho,\sigma}(P)\ne(0,0)\ne\mathrm{st}_{\rho,\sigma}(Q), &&\mathrm{st}_{\rho,\sigma}(P)\sim\mathrm{st}_{\rho,\sigma}(Q),\\
&\mathrm{en}_{\rho,\sigma}(P)\ne(0,0)\ne\mathrm{en}_{\rho,\sigma}(Q), &&\mathrm{en}_{\rho,\sigma}(P)\sim\mathrm{en}_{\rho,\sigma}(Q).
\end{aligned}$$

4 "Fixed points" of (ρ,σ)-brackets

The aim of this section is to construct $F\in W$ such that $[F,P]_{\rho,\sigma}=\ell_{\rho,\sigma}(P)$ and $[F,Q]_{\rho,\sigma}=\ell_{\rho,\sigma}(Q)$ for irreducible pairs (P,Q) and some given $(\rho,\sigma)\in\mathfrak{V}$.

Lemma 4.1. *Let $m,n\in\mathbb{N}$. If $\gcd(n,m)=1$, then for all $r>mn-n-m$ there exist $a,b\ge0$ such that $na+mb=r$.*

Proof. There exist $\bar a,\bar b\in\mathbb{Z}$ such that $r+n+m=\bar an+\bar bm$. We can assume that $0<\bar a\le m$. Then $\bar an\le mn<r+n+m$, and so $\bar b>0$. Hence $a:=\bar a-1$ and $b:=\bar b-1$ satisfy the thesis. □

Lemma 4.2. *Let $m,n,r\in\mathbb{N}$. If $c=\gcd(n,m)$ and $\bar c=\gcd(c,r)$, then there exist $a,b\ge0$ such that $\bar c=\gcd(an+bm,r)$.*

Proof. There exist $a,\bar{b}\ge 0$ and $\alpha,\beta>0$ such that $c=an-\bar{b}m$ and $\bar{c}=\alpha r-\beta c$. Let k be such that $kr>\bar{b}$ and set $b:=kr-\bar{b}>0$. A direct computation shows that

$$\bar{c}=r(\alpha+m\beta k)-(an+bm)\beta,$$

Hence $\gcd(an+bm,r)$ divides $\bar{c}$. But, since $\bar{c}=\gcd(c,r)$ and $c=\gcd(m,n)$, we have that $\bar{c}$ divides $an+bm$ and r, and so $\bar{c}=\gcd(an+bm,r)$. □

Theorem 4.3. *Let* $(\rho,\sigma)\in\mathfrak{V}$ *and* $C\in W$ *such that*

$$v_{\rho,\sigma}(C)>0 \quad \text{and} \quad \ell_{\rho,\sigma}(C)\ne\zeta\ell_{\rho,\sigma}(D^h), \quad \text{for all } D\in W,\ \zeta\in K^{\times} \text{ and } h>1.$$

If there exist $n,m\in\mathbb{N}$ *and* $A,B\in W$ *such that*

1. $\ell_{\rho,\sigma}(A)=\ell_{\rho,\sigma}(C^n)$,
2. $\ell_{\rho,\sigma}(B)=\ell_{\rho,\sigma}(C^m)$,
3. $c:=\gcd(n,m)\notin\{n,m\}$,
4. $\ell_{\rho,\sigma}([A,B])=\lambda\ell_{\rho,\sigma}(C^h)$, *for some* $h\in\mathbb{N}_0$ *and* $\lambda\in K^{\times}$,

then there exist $D\in W$, $\mu\in K^{\times}$ *and* $k,j_0\in\mathbb{N}$, *such that*

$$[D,C^k]_{\rho,\sigma}=\mu\ell_{\rho,\sigma}(C^{k+j_0}).$$

Proof. Take A and B satisfying the hypothesis of the statement with c minimum. Set

$$m_1:=m/c,\quad n_1:=n/c,\quad D_0:=A^{m_1}-B^{n_1}$$

and

$$X:=\left\{D=D_0+\sum_{i,j\in\mathbb{N}_0}\lambda_{ij}A^iB^j\in W: in+jm<cn_1m_1 \text{ and } \lambda_{ij}\in K\right\}.$$

We claim that each $D\in X$ satisfies

$$\ell_{\rho,\sigma}([D,B])=m_1\lambda\ell_{\rho,\sigma}(C^{nm_1-n+h}) \tag{4.1}$$

and

$$\ell_{\rho,\sigma}([D,A])=n_1\lambda\ell_{\rho,\sigma}(C^{mn_1-m+h}). \tag{4.2}$$

In fact, this is true for D_0 since, by Proposition 1.9, Lemma 1.25, and items (1) and (4), we have

$$\ell_{\rho,\sigma}([D_0,B])=\ell_{\rho,\sigma}([A^{m_1},B])=m_1\lambda\ell_{\rho,\sigma}(C^{nm_1-n+h}).$$

and similarly

$$\ell_{\rho,\sigma}([D_0,A])=n_1\lambda\ell_{\rho,\sigma}(C^{mn_1-m+h}).$$

In particular $D_0\ne 0$. So, in order to establish (4.1) and (4.2), it suffices to show that

$$v_{\rho,\sigma}([A^iB^j,B])<(nm_1-n+h)v_{\rho,\sigma}(C)$$

and

$$v_{\rho,\sigma}([A^iB^j,A])<(mn_1-m+h)v_{\rho,\sigma}(C),$$

for all i,j such that $in+jm<n_1m_1c$. But this follows from the fact that, again by Proposition 1.9, Lemma 1.25, and items (1), (2) and (4),

$$\ell_{\rho,\sigma}([A^iB^j,B])=\ell_{\rho,\sigma}([A^i,B]B^j)=i\lambda\ell_{\rho,\sigma}(C^{ni+mj-n+h})$$

and

$$\ell_{\rho,\sigma}([A^iB^j,A])=\ell_{\rho,\sigma}(A^i[B^j,A])=j\lambda\ell_{\rho,\sigma}(C^{ni+mj-m+h}).$$

Now, by Remark 1.13, equality (4.1) implies that for $D\in X$

$$v_{\rho,\sigma}(D)+v_{\rho,\sigma}(B)-(\rho+\sigma)\ge v_{\rho,\sigma}(C^{nm_1-n+h}),$$

and so, by item (2),

$$v_{\rho,\sigma}(D) > v_{\rho,\sigma}(C^{nm_1-n+h}) - v_{\rho,\sigma}(B) = (nm_1 - m - n + h)v_{\rho,\sigma}(C) \tag{4.3}$$

for all $D \in X$. Hence there exists $D_1 \in X$ such that $v_{\rho,\sigma}(D_1)$ is minimum. We have two alternatives:

$$[D_1, B]_{\rho,\sigma} \neq 0 \quad \text{or} \quad [D_1, B]_{\rho,\sigma} = 0. \tag{4.4}$$

Note that

$$j_0 := nm_1 - n - m + h \geq c(n_1m_1 - n_1 - m_1) = c\big((n_1-1)(m_1-1)-1\big) > 0,$$

since $\gcd(n_1, m_1) = 1$ and $n_1, m_1 > 1$ by item (3). Hence, in the first case, the thesis holds with $k = m$ and $\mu = m_1\lambda$, because, by item (2), Corollary 1.19 and (4.1),

$$[D_1, C^m]_{\rho,\sigma} = [D_1, B]_{\rho,\sigma} = m_1\lambda\ell_{\rho,\sigma}(C^{nm_1-n+h}).$$

Assume now that $[D_1, B]_{\rho,\sigma} = 0$. We are going to show that this alternative is impossible, because it implies that c is not minimum. In other words, that

(*) there exist $\bar{A}, \bar{B} \in W$, $\bar{\lambda} \in K^\times$ and $\bar{n}, \bar{m}, \bar{c}, \bar{h} \in \mathbb{N}$ with $\bar{c} < c$, such that (1), (2), (3) and (4) hold, with $\bar{A}$, $\bar{B}$, $\bar{\lambda}$, $\bar{n}$, $\bar{m}$, $\bar{c}$ and $\bar{h}$ instead of A, B, λ, n, m, c and h respectively.

With this purpose in mind, we claim that there exist $\lambda_1 \in K^\times$ and $r \in \mathbb{N}$, such that

$$\ell_{\rho,\sigma}(D_1) = \lambda_1\ell_{\rho,\sigma}(C^r), \quad r < n_1m_1c, \quad r > c \quad \text{and} \quad c \nmid r \tag{4.5}$$

In fact, by Corollary 1.19 and item (2), we know that $[D_1, C^m]_{\rho,\sigma} = [D_1, B]_{\rho,\sigma} = 0$, which by Lemma 1.25 implies that $[D_1, C]_{\rho,\sigma} = 0$. Hence, by item (2) of Theorem 1.22, there exists $R = \ell_{\rho,\sigma}(R) \in L$, $\zeta, \xi \in K^\times$ and $r, s \in \mathbb{N}$, such that

$$\ell_{\rho,\sigma}(D_1) = \zeta R^r \quad \text{and} \quad \ell_{\rho,\sigma}(C) = \xi R^s.$$

Besides, by the conditions required to C, it must be $s = 1$ and so, by Proposition 1.9, $\ell_{\rho,\sigma}(D_1) = \frac{\zeta}{\xi^r}\ell_{\rho,\sigma}(C^r)$, which proves the equality in (4.5) with $\lambda_1 := \frac{\zeta}{\xi^r}$. Moreover

$$rv_{\rho,\sigma}(C) = v_{\rho,\sigma}(D_1) \leq v_{\rho,\sigma}(D_0) = v_{\rho,\sigma}(A^{m_1} - B^{n_1}) < n_1m_1cv_{\rho,\sigma}(C),$$

where the last inequality follows from the fact that, by items (1) and (2),

$$\ell_{\rho,\sigma}(A^{m_1}) = \ell_{\rho,\sigma}(C^{cn_1m_1}) = \ell_{\rho,\sigma}(B^{n_1}).$$

Thus $r < n_1m_1c$. Note that by (4.3) and the equality in (4.5),

$$rv_{\rho,\sigma}(C) = v_{\rho,\sigma}(D_1) > (nm_1 - m - n + h)v_{\rho,\sigma}(C) \geq c(m_1n_1 - m_1 - n_1)v_{\rho,\sigma}(C).$$

Hence

$$r > c(m_1n_1 - m_1 - n_1) = c\big((m_1-1)(n_1-1)-1\big) \geq c, \tag{4.6}$$

where the last inequality holds, as before, because $m_1, n_1 \geq 2$ and $m_1 \neq n_1$. Next we will prove that c does not divide r. Assume on the contrary that $c \mid r$. By Lemma 4.1 and the first inequality in (4.6) there exist $a_1, b_1 \geq 0$ such that

$$a_1n_1 + b_1m_1 = \frac{r}{c}.$$

Consequently

$$a_1n + b_1m = r < cn_1m_1,$$

and so $D_2 := D_1 - \lambda_1A^{a_1}B^{b_1} \in X$. Moreover, since by items (1) and (2), and the equality in (4.5),

$$\lambda_1\ell_{\rho,\sigma}(A^{a_1}B^{b_1}) = \lambda_1\ell_{\rho,\sigma}(C^{a_1n+b_1m}) = \lambda_1\ell_{\rho,\sigma}(C^r) = \ell_{\rho,\sigma}(D_1),$$

we get $v_{\rho,\sigma}(D_2) < v_{\rho,\sigma}(D_1)$, which contradicts the minimality of $v_{\rho,\sigma}(D_1)$. Thus c does not divide r.

Set $\bar{c} := \gcd(c,r)$ and $\bar{A} := \frac{1}{\lambda_1} D_1$. By (4.5), we know that

$$\ell_{\rho,\sigma}(\bar{A}) = \ell_{\rho,\sigma}(C^{\bar{n}}),$$

where $\bar{n} := r > c > \bar{c}$. Moreover, by Lemma 4.2 there exist $a, b \geq 0$, such that $\gcd(r, an + bm) = \bar{c}$. Note that $a > 0$ or $b > 0$, because $\bar{c} \neq r$. In particular $\bar{c} < c \leq \min(n,m) \leq an + bm$. Let $\bar{B} := A^a B^b$. By Proposition 1.9 and items (1) and (2),

$$\ell_{\rho,\sigma}(\bar{B}) = \ell_{\rho,\sigma}(C^{\bar{m}}),$$

where $\bar{m} := an + bm$. So, in order to verify that (*) holds, it only remains to establish (4). But, since,

$$\lambda_1 [\bar{A}, \bar{B}] = [D_1, A^a B^b] = [D_1, A^a] B^b + A^a [D_1, B^b].$$

and, by Proposition 1.9, Lemma 1.25, items (1) and (2), and equalities (4.1) and (4.2),

$$\ell_{\rho,\sigma}([D_1, A^a] B^b) = a n_1 \lambda \ell_{\rho,\sigma}(C^{\bar{m} + c(m_1 n_1 - m_1 - n_1) + h})$$

and

$$\ell_{\rho,\sigma}(A^a [D_1, B^b]) = b m_1 \lambda \ell_{\rho,\sigma}(C^{\bar{m} + c(m_1 n_1 - m_1 - n_1) + h}),$$

we have

$$\ell_{\rho,\sigma}([\bar{A}, \bar{B}]) = \bar{\lambda} C^{\bar{h}},$$

with

$$\bar{\lambda} := \frac{\lambda}{\lambda_1}(a n_1 + b m_1) \neq 0 \quad \text{and} \quad \bar{h} := \bar{m} + c(m_1 n_1 - m_1 - n_1) + h > 0,$$

as desired. □

Corollary 4.4. *Let* $(\rho,\sigma) \in \mathfrak{V}$ *and let* $C \in W$ *such that*

$$v_{\rho,\sigma}(C) > 0 \quad \text{and} \quad \ell_{\rho,\sigma}(C) \neq \zeta \ell_{\rho,\sigma}(D^h) \quad \text{for all } D \in W,\ \zeta \in K^{\times} \text{ and } h \in \mathbb{N}.$$

If there exist $n, m \in \mathbb{N}$ *and* $A, B \in W$ *such that*

1. $\ell_{\rho,\sigma}(A) = \ell_{\rho,\sigma}(C^n)$,
2. $\ell_{\rho,\sigma}(B) = \ell_{\rho,\sigma}(C^m)$,
3. $c := \gcd(n,m) \notin \{n, m\}$,
4. $\ell_{\rho,\sigma}([A,B]) = \lambda \ell_{\rho,\sigma}(C^h)$, *for some* $h \in \mathbb{N}_0$ *and* $\lambda \in K^{\times}$,

then

$$\operatorname{Supp}(\ell_{\rho,\sigma}(C)) \neq \{(j,j)\} \quad \text{for all } j.$$

Proof. Assume on the contrary that $\ell_{\rho,\sigma}(C) = d x^j y^j$ for some $d \in K^{\times}$ and some $j \in \mathbb{N}_0$. By Theorem 4.3 there exist $D \in W$, $\mu \in K^{\times}$, $k \in \mathbb{N}$ and $j_0 \in \mathbb{N}_0$, such that $[D, C^k]_{\rho,\sigma} = \mu C^{k+j_0}$. By Corollary 1.19, we can assume that D is (ρ,σ)-homogeneous. Write $D = \sum d_{rs} X^r Y^s$. Since, by Remark 1.13,

$$[d_{rr} X^r Y^r, d^k X^{kj} Y^{kj}] = 0$$

and, by Lemma 1.7,

$$\ell_{\rho,\sigma}([d_{rs} X^r Y^s, d^k X^{kj} Y^{kj}]) = (s-r) kj d_{rs} d^k x^{r+kj-1} y^{s+kj-1} \quad \text{for all } r \neq s,$$

we have

$$\ell_{\rho,\sigma}([D, d^k X^{kj} Y^{kj}]) = \sum_{r \neq s} (s-r) kj d_{rs} d^k x^{r+kj-1} y^{s+kj-1} \tag{4.7}$$

On the other hand, by Proposition 1.9 and Corollary 1.19,

$$\ell_{\rho,\sigma}([D, d^k X^{kj} Y^{kj}]) = \ell_{\rho,\sigma}([D, C^k]) = \mu \ell_{\rho,\sigma}(C^{k+j_0}) = \mu d^{k+j_0} x^{(k+j_0)j} y^{(k+j_0)j},$$

which contradicts (4.7). □

Theorem 4.5. *Let* (P,Q) *be an irreducible pair and let* $(\rho,\sigma) \in \mathfrak{V}$.

(1) *If* $(\rho,\sigma)\notin \mathrm{Val}(P)$, *then* $v_{1,-1}(\ell_{\rho,\sigma}(P))\neq 0$.
(2) $v_{1,-1}(\mathrm{st}_{\rho,\sigma}(P))\neq 0$,
(3) *There exists a* (ρ,σ)*-homogeneous element* $F\in W$, *such that*

$$[P,F]_{\rho,\sigma}=\ell_{\rho,\sigma}(P)\quad\text{and}\quad [Q,F]_{\rho,\sigma}=\xi\ell_{\rho,\sigma}(Q),$$

where $\xi=v_{\rho,\sigma}(Q)/v_{\rho,\sigma}(P)$. *Moreover* $v_{\rho,\sigma}(F)=\rho+\sigma$.


Y
Q
P
C
F
1
1
X
$[F,P]_{2,-1}=\ell_{2,-1}(P)$
$[F,Q]_{2,-1}=\frac{3}{2}\ell_{2,-1}(Q)$

FIGURE 14. Theorem 4.5.

Proof. By Proposition 3.10, there exist $C\in W$, $\lambda_P,\lambda_Q\in K^{\times}$ and $n,m\in\mathbb{N}$ such that

(1) $v_{\rho,\sigma}(C)>0$ and $\ell_{\rho,\sigma}(C)\neq\zeta\ell_{\rho,\sigma}(D^h)$ for all $D\in W$, $\zeta\in K^{\times}$ and $h>1$,
(2) $\ell_{\rho,\sigma}(P)=\lambda_P\ell_{\rho,\sigma}(C^m)$,
(3) $\ell_{\rho,\sigma}(Q)=\lambda_Q\ell_{\rho,\sigma}(C^n)$,
(4) $\gcd(n,m)\notin\{n,m\}$.

Thus, the conditions of Theorem 4.3 and Corollary 4.4 are fulfilled with

$$A:=\frac{1}{\lambda_Q}Q,\quad B:=\frac{1}{\lambda_P}P,\quad h:=0\quad\text{and}\quad \lambda:=\frac{1}{\lambda_P\lambda_Q}.$$

In particular, if $(\rho,\sigma)\notin\mathrm{Val}(P)$, then $\mathrm{Supp}(\ell_{\rho,\sigma}(C))=\{(i,j)\}$ with $i\neq j$, and so, $\mathrm{Supp}(\ell_{\rho,\sigma}(P))=\{(mi,mj)\}$. Item (1) follows immediately from this fact. In order to prove item (2), we note that, by item (3) of Proposition 2.8 there exist $(\rho',\sigma')<(\rho,\sigma)$ such that $\mathrm{Supp}(\ell_{\rho',\sigma'}(P))=\{\mathrm{st}_{\rho,\sigma}(P)\}$ and so, by item (1)

$$v_{1,-1}(\mathrm{st}_{\rho,\sigma}(P))=v_{1,-1}(\ell_{\rho',\sigma'}(P))\neq 1$$

We now prove item (3). By Theorem 4.3, there exist $D\in W$, $\mu\in K^{\times}$ and $k,j_0\in\mathbb{N}$, such that

$$[D,C^k]_{\rho,\sigma}=\mu C^{k+j_0},$$

and so, by Theorem 1.26, there exists a (ρ,σ)-homogeneous element $E\in W$, such that

$$[C^t,E]_{\rho,\sigma}=t\ell_{\rho,\sigma}(C^t)\quad\text{for all } t\in\mathbb{N}.$$

Hence, by Corollary 1.19,

$$[P,E]_{\rho,\sigma}=[\lambda_P C^m,E]_{\rho,\sigma}=m\ell_{\rho,\sigma}(\lambda_P C^m)=m\ell_{\rho,\sigma}(P)$$

and

$$[Q,E]_{\rho,\sigma}=[\lambda_Q C^n,E]_{\rho,\sigma}=n\ell_{\rho,\sigma}(\lambda_Q C^n)=n\ell_{\rho,\sigma}(Q).$$

If we set $F:=\frac{1}{m}E$, then we have

$$[P,F]_{\rho,\sigma}=\ell_{\rho,\sigma}(P)\quad\text{and}\quad [Q,F]_{\rho,\sigma}=\frac{n}{m}\ell_{\rho,\sigma}(Q).$$

Note now that $v_{\rho,\sigma}(P)=mv_{\rho,\sigma}(C)$ and $v_{\rho,\sigma}(Q)=nv_{\rho,\sigma}(C)$, and so

$$\xi:=\frac{n}{m}=\frac{v_{\rho,\sigma}(Q)}{v_{\rho,\sigma}(P)}.$$

Finally, it is clear that $[P,F]_{\rho,\sigma}=\ell_{\rho,\sigma}(P)$ implies $v_{\rho,\sigma}(F)=\rho+\sigma$. □

Proposition 4.6. *Let $(\rho,\sigma)\in\mathfrak{V}$, let $P,F\in W\setminus\{0\}$ and let $f_{F,\rho,\sigma}$ and $f_{P,\rho,\sigma}$ be as in Definition 1.20. Assume that F is (ρ,σ)-homogeneous and $[P,F]_{\rho,\sigma}=\ell_{\rho,\sigma}(P)$. Then $f_{F,\rho,\sigma}$ is separable and every irreducible factor of $f_{P,\rho,\sigma}$ divides $f_{F,\rho,\sigma}$.*

Proof. By item (1) of Theorem 1.22, there exist $h\ge 0$ and $c\in\mathbb{Z}$, such that

$$x^h f_P=cf_Pf_F+axf_P'f_F-bxf_F'f_P,$$

where $a:=v_{\rho,\sigma}(F)$, $b:=v_{\rho,\sigma}(P)$, $f_P:=f_{P,\rho,\sigma}$ and $f_F:=f_{F,\rho,\sigma}$. Hence, the pair of polynomials (f_P,f_F) satisfies $\mathrm{PE}(1,0,a,b,c)$. Since $f_P\ne 0\ne f_F$ the result follows from Proposition 1.24. □

5 Irreducible pairs and subrectangular elements

In this section we prove that each irreducible pair can be transformed into an irreducible pair whose elements are subrectangular.

Lemma 5.1. *Let $A_i\in W\setminus\{0\}$ $(i=0,\dots,n)$ and let $(\rho,\sigma)\in\mathfrak{V}$. Suppose that there exists $q\in\mathbb{Q}$ such that $v_{\rho,\sigma}(A_i)=q$ for all i and set $A:=\sum_{i=0}^n A_i$. Then*

$$A\ne 0\ \textit{and}\ v_{\rho,\sigma}(A)=q\iff\sum_i \ell_{\rho,\sigma}(A_i)\ne 0$$
$$\iff A\ne 0\ \textit{and}\ \ell_{\rho,\sigma}(A)=\sum_i\ell_{\rho,\sigma}(A_i).$$

Proof. This is clear since the isomorphism of K-vector spaces $\Psi\colon W\to L$, introduced at the beginning of Section 1, preserves the (ρ,σ)-degree. □

For $\varphi\in\mathrm{Aut}(W)$, we will denote by φ_L the automorphism of L defined by $\varphi_L(x):=\Psi(\varphi(X))$ and $\varphi_L(y):=\Psi(\varphi(Y))$.

Proposition 5.2. *Let $(\rho,\sigma)\in\mathfrak{V}$, $\lambda\in K$ and φ an automorphism of W. In any of the following cases:*

1. *$P\in W\setminus\{0\}$, $(\rho,\sigma)=(\rho,1)$, $\varphi(X)=X+\lambda Y^\rho$ and $\varphi(Y)=Y$,*
2. *$P\in W\setminus\{0\}$, $(\rho,\sigma)=(1,\sigma)$, $\varphi(X)=X$ and $\varphi(Y)=Y+\lambda X^\sigma$,*

we have

$$\ell_{\rho,\sigma}(\varphi(P))=\varphi_L(\ell_{\rho,\sigma}(P))\quad\textit{and}\quad v_{\rho,\sigma}(\varphi(P))=v_{\rho,\sigma}(P).$$

Furthermore, in the case (1),

$$\ell_{\rho_1,\sigma_1}(\varphi(P))=\ell_{\rho_1,\sigma_1}(P)\quad\textit{for all } (1,-1)<(\rho_1,\sigma_1)<(\rho,\sigma),$$

and, in the case (2),

$$\ell_{\rho_1,\sigma_1}(\varphi(P))=\ell_{\rho_1,\sigma_1}(P)\quad\textit{for all } (\rho,\sigma)<(\rho_1,\sigma_1)<(-1,1).$$

Proof. We prove (1) and leave (2), which are similar, to the reader. By item (3) of Proposition 1.9,

$$v_{\rho,1}\bigl(\varphi(X^iY^j)\bigr) = iv_{\rho,1}(X+\lambda Y^{\rho})+jv_{\rho,1}(Y) = \rho i+j = v_{\rho,1}(X^iY^j) \quad \text{for all } i,j\in\mathbb{N}_0.$$

Hence

$$v_{\rho,1}(\varphi(P)) = v_{\rho,1}(P) \quad \text{for all } P\in W\setminus\{0\}, \tag{5.1}$$

since φ is bijective and therefore induce and isomorphism between the gradations associate with the (ρ,σ)-filtrations. We fix now a $P\in W\setminus\{0\}$ and write

$$P = \sum_{i=0}^{n} \lambda_i X^{r-i}Y^{s+i\rho} + R,$$

where $R=0$ or $v_{\rho,1}(R) < v_{\rho,1}(P)$. By (5.1), Lemma 5.1 and item (2) of Proposition 1.9,

$$\begin{aligned}
\ell_{\rho,1}(\varphi(P)) &= \ell_{\rho,1}\left(\sum_{i=0}^{n}\lambda_i\varphi\bigl(X^{r-i}Y^{s+i\rho}\bigr)\right)\\
&= \ell_{\rho,1}\left(\sum_{i=0}^{n}\lambda_i(X+\lambda Y^{\rho})^{r-i}Y^{s+i\rho}\right)\\
&= \sum_{i=0}^{n}\lambda_i\ell_{\rho,1}\bigl((X+\lambda Y^{\rho})^{r-i}Y^{s+i\rho}\bigr)\\
&= \sum_{i=0}^{n}\lambda_i(x+\lambda y^{\rho})^{r-i}y^{s+i\rho}\\
&= \varphi_L\left(\sum_{i=0}^{n}\lambda_i x^{r-i}y^{s+i\rho}\right)\\
&= \varphi_L(\ell_{\rho,1}(P)),
\end{aligned}$$

as desired. Let $(\rho_1,\sigma_1)\in\mathfrak{V}$ such that $(\rho_1,\sigma_1)<(\rho,1)$. Then $\rho_1>\rho\sigma_1$, and so

$$\ell_{\rho_1,\sigma_1}(X+\lambda Y^{\rho}) = x.$$

Hence, by Proposition 1.9,

$$\ell_{\rho_1,\sigma_1}(\varphi(X^iY^j)) = \ell_{\rho_1,\sigma_1}((X+\lambda Y^{\rho})^iY^j) = x^iy^j$$

and

$$v_{\rho_1,\sigma_1}(\varphi(X^iY^j)) = i\rho_1 + j\sigma_1 = v_{\rho_1,\sigma_1}(X^iY^j),$$

which implies that

$$v_{\rho_1,\sigma_1}(\varphi(R)) = v_{\rho_1,\sigma_1}(R) \quad \text{for all } R\in W\setminus\{0\}. \tag{5.2}$$

Fix now $P\in W\setminus\{0\}$ and write

$$P = \sum_{\{(i,j):\rho_1 i+\sigma_1 j = v_{\rho_1,\sigma_1}(P)\}} \lambda_{ij}X^iY^j + R,$$

with $R=0$ or $v_{\rho_1,\sigma_1}(R)<v_{\rho_1,\sigma_1}(P)$. Again by (5.1), Lemma 5.1 and item (2) of Proposition 1.9

$$\begin{aligned}\ell_{\rho_1,\sigma_1}(\varphi(P)) &= \ell_{\rho_1,\sigma_1}\Bigl(\sum_{i,j}\lambda_{ij}\varphi(X^iY^j)\Bigr)\\ &= \ell_{\rho_1,\sigma_1}\Bigl(\sum_{i,j}\lambda_{ij}(X+\lambda Y^{\rho})^iY^j\Bigr)\\ &= \sum_{i,j}\lambda_{ij}\ell_{\rho_1,\sigma_1}\bigl((X+\lambda Y^{\rho})^iY^j\bigr)\\ &= \sum_{i,j}\lambda_{ij}x^iy^j\\ &= \ell_{\rho_1,\sigma_1}(P),\end{aligned}$$

as desired. □

We let $\mathrm{CH}(D)$ denote the convex hull of a subset D of $\mathbb{R}^2$.

Remark 5.3. Let $(\rho,\sigma)\in\mathfrak{V}$ and let $A\in W\setminus\{0\}$. Note that if

$$\ell_{\rho,\sigma}(A)=\sum_{i=0}^{\gamma}a_ix^{r-i\sigma}y^{s+i\rho}\quad\text{with } a_0\neq 0 \text{ and } a_\gamma\neq 0,$$

then

$$\mathrm{CH}(\mathrm{Supp}(\ell_{\rho,\sigma}(A)))=\{\lambda(r,s)+(1-\lambda)(r-\gamma\sigma,s+\gamma\rho):0\le\lambda\le 1\}.$$

Note that by Proposition 1.9,

$$\mathrm{CH}(\mathrm{Supp}(\ell_{\rho,\sigma}(A^t)))=\{\lambda t(r,s)+(1-\lambda)t(r-\gamma\sigma,s+\gamma\rho):0\le\lambda\le 1\}. \tag{5.3}$$

for all $t\in\mathbb{N}$.

Remark 5.4. Let $(\rho,\sigma)\in\mathfrak{V}$ and $P,F\in W\setminus\{0\}$. Assume that F is (ρ,σ)-homogeneous and that $[P,F]_{\rho,\sigma}=\ell_{\rho,\sigma}(P)$. Since $v_{\rho,\sigma}(F)=\rho+\sigma$,

$$\mathrm{Supp}(F)\subseteq R_{\rho,\sigma}=\{(x,y)\in\mathbb{R}^2:x\rho+y\sigma=\rho+\sigma\}, \tag{5.4}$$

where $R_{\rho,\sigma}:=\{(1,1)+t(-\sigma,\rho):t\in\mathbb{R}\}$.

Proposition 5.5. *Let P, F and (ρ,σ) be as in Remark 5.4. Set*

$$\mathcal{P}:=\mathrm{CH}(\mathrm{Supp}(\ell_{\rho,\sigma}(P)))\quad\textit{and}\quad\mathrm{Cone}(\mathcal{P}):=\{tu:t>0,\ t\in\mathbb{R}\ \textit{and}\ u\in\mathcal{P}\}.$$

We have

$$\{\mathrm{st}_{\rho,\sigma}(F),\mathrm{en}_{\rho,\sigma}(F)\}\subseteq(\mathrm{Cone}(\mathcal{P})\cap R_{\rho,\sigma}\cap\mathbb{N}_0^2)\cup\{(1,1)\},$$

where $R_{\rho,\sigma}$ is as above.

Proof. For the sake of brevity we will write $\mathrm{st}(F)$ and $\mathrm{en}(F)$ instead of $\mathrm{st}_{\rho,\sigma}(F)$ and $\mathrm{en}_{\rho,\sigma}(F)$, respectively. By Remark 5.4, we know that $\mathrm{Supp}(F)\subseteq R_{\rho,\sigma}$, and so $\{\mathrm{st}(F),\mathrm{en}(F)\}\subseteq R_{\rho,\sigma}\cap\mathbb{N}_0^2$. We claim that if one of $\mathrm{st}(F)$ or $\mathrm{en}(F)$ is not in $\mathrm{Cone}(\mathcal{P})$, then it is equal to $(1,1)$. For this, note that $\mathrm{st}(F)\sim\mathrm{st}(P)$ implies $\mathrm{st}(F)\in\mathrm{Cone}(\mathcal{P})$. Consequently, from item (1) of Proposition 1.17 and the fact that $[P,F]_{\rho,\sigma}=\ell_{\rho,\sigma}(P)$ it follows that if $\mathrm{st}(F)\notin\mathrm{Cone}(\mathcal{P})$, then $\mathrm{st}(F)=(1,1)$. A similar argument works for $\mathrm{en}(F)$. □

Lemma 5.6. *Let P, F and (ρ,σ) be as in Remark 5.4. If F is a monomial, then $(\rho,\sigma)\notin\mathrm{Val}(P)$ and $F=\mu XY$ for some $\mu\in K^{\times}$.*



Figure 15. Proposition 5.5.

Proof. Write

$$\ell_{\rho,\sigma}(P) = \sum_{i=0}^{\alpha} \lambda_i x^{r-i\sigma} y^{s+i\rho} \quad \text{with } \lambda_0, \lambda_\alpha \neq 0.$$

By Definition 1.15, the equality $[P, F]_{\rho,\sigma} = \ell_{\rho,\sigma}(P)$ implies $v_{\rho,\sigma}(F) = \rho + \sigma$. Hence there exists $j \in \mathbb{Z}$ such that

$$F = \mu X^{1-j\sigma} Y^{1+j\rho}.$$

Now, by Proposition 1.18,

$$[P, F]_{\rho,\sigma} = \sum_{i=0}^{\alpha} \mu \lambda_i \mu_i x^{r-(i+j)\sigma} y^{s+(i+j)\rho},$$

where

$$\mu_i = (1 - j\sigma)(s + i\rho) - (1 + j\rho)(r - i\sigma) = (s - r) - j(r\rho + s\sigma) + i(\rho + \sigma). \quad (5.5)$$

Consequently,

$$\sum_{i=0}^{\alpha} \mu \lambda_i \mu_i x^{r-(i+j)\sigma} y^{s+(i+j)\rho} = \sum_{i=0}^{\alpha} \lambda_i x^{r-i\sigma} y^{s+i\rho} \neq 0. \quad (5.6)$$

Since

$$v_{1,-1}\left(\sum_{i=0}^{\alpha} \lambda_i x^{r-i\sigma} y^{s+i\rho}\right) = r - s$$

and

$$v_{1,-1}\left(\sum_{i=0}^{\alpha} \mu \lambda_i \mu_i x^{r-(i+j)\sigma} y^{s+(i+j)\rho}\right) \leq r - s - j(\rho + \sigma),$$

from equality (5.6), it follows that necessarily $j \leq 0$. Similarly, since

$$v_{-1,1}\left(\sum_{i=0}^{\alpha} \lambda_i x^{r-i\sigma} y^{s+i\rho}\right) = s - r + \alpha(\rho + \sigma)$$

and

$$v_{-1,1}\left(\sum_{i=0}^{\alpha} \mu \lambda_i \mu_i x^{r-(i+j)\sigma} y^{s+(i+j)\rho}\right) \leq s - r + (\alpha + j)(\rho + \sigma),$$

necessarily $j \geq 0$. Hence $F = \mu XY$ and equations (5.5) and (5.6) become

$$\mu_i = (r - s) - i(\rho + \sigma) \tag{5.7}$$

and

$$\sum_{i=0}^{\alpha} \mu \lambda_i \mu_i x^{r-i\sigma} y^{s+i\rho} = \sum_{i=0}^{\alpha} \lambda_i x^{r-i\sigma} y^{s+i\rho}. \tag{5.8}$$

Consequently, $\mu_0 = \mu_\alpha$, which, by (5.7), implies that $\alpha = 0$. ☐

Proposition 5.7. *Let P, F and (ρ, σ) be as in Remark 5.4. We have:*

1. *If $(\rho, \sigma) = (1, 1)$, then $\operatorname{Supp}(F) \subseteq \{(2, 0), (1, 1), (0, 2)\}$. Moreover,*
    - *if $(0, 2) \in \operatorname{Supp}(F)$, then $(0, l) \in \operatorname{Supp}(\ell_{\rho,\sigma}(P))$ for some $l \in \mathbb{N}$,*
    - *if $(2, 0) \in \operatorname{Supp}(F)$, then $(l, 0) \in \operatorname{Supp}(\ell_{\rho,\sigma}(P))$ for some $l \in \mathbb{N}$.*
2. *If $\rho > \sigma > 0$ and $(\rho, \sigma) \in \operatorname{Val}(P)$, then $\sigma = 1$, $\operatorname{Supp}(F) = \{(1, 1), (0, 1+\rho)\}$ and $(0, l) \in \operatorname{Supp}(\ell_{\rho,\sigma}(P))$ for some $l \in \mathbb{N}$.*
3. *If $\sigma > \rho > 0$ and $(\rho, \sigma) \in \operatorname{Val}(P)$, then $\rho = 1$, $\operatorname{Supp}(F) = \{(1, 1), (1+\sigma, 0)\}$ and $(l, 0) \in \operatorname{Supp}(\ell_{\rho,\sigma}(P))$ for some $l \in \mathbb{N}$.*

Consequently, the polynomial $f_{F,\rho,\sigma}$, introduced in Definition 1.20, satisfies

- $\deg(f_{F,1,1}) \leq 2$,
- $\deg(f_{F,1,1}) = 2$ *if and only if* $\{(2, 0), (0, 2)\} \subseteq \operatorname{Supp}(F)$,
- *if $\sigma > \rho > 0$ or $\rho > \sigma > 0$, then* $\deg(f_{F,\rho,\sigma}) = 1$.



(A) Case $\rho = \sigma = 1$ (B) Case $\rho > \sigma > 0$

FIGURE 16. Proposition 5.7.

Proof. Note that if $(1, 1) + t(-\sigma, \rho) \in \mathbb{N}_0^2$, then $t \in \mathbb{Z}$, $1 - \sigma t \geq 0$ and $1 + t\rho \geq 0$. Since $\sigma, \rho \geq 1$ we obtain that

$$-1 \leq -\frac{1}{\rho} \leq t \leq \frac{1}{\sigma} \leq 1. \tag{5.9}$$

Hence

$$(1, 1) + t(-\sigma, \rho) \in \mathbb{N}_0^2 \Longrightarrow \begin{cases} t \in \{-1, 0, 1\} & \text{if } \rho = \sigma = 1, \\ t \in \{0, 1\} & \text{if } \rho > \sigma \geq 1, \\ t \in \{-1, 0\} & \text{if } \sigma > \rho \geq 1. \end{cases}$$

Consequently, by Remark 5.4,

$$\operatorname{Supp}(F) \subseteq \begin{cases} \{(2,0),(1,1),(0,2)\} & \text{if } \rho=\sigma=1,\\ \{(1,1),(1-\sigma,1+\rho)\} & \text{if } \rho>\sigma\geq 1,\\ \{(1,1),(1+\sigma,1-\rho)\} & \text{if } \sigma>\rho\geq 1.\end{cases}$$

This finishes the proof of the first part of item (1). Moreover, by Proposition 5.5, we know that if $(0,2) \in \operatorname{Supp}(F)$, then $(0,2) \in \operatorname{Cone}(\mathcal{P})$, and so

$$(0,l) \in \operatorname{Supp}(\ell_{\rho,\sigma}(P)) \quad \text{for some } l \in \mathbb{N},$$

and similarly if $(2,0) \in \operatorname{Supp}(F)$, then $(l,0) \in \operatorname{Supp}(\ell_{\rho,\sigma}(P))$ for some $l \in \mathbb{N}$. Now we prove item (2). By Lemma 5.6, if $(\rho,\sigma) \in \operatorname{Val}(P)$ then F is not a monomial, and so $\sigma = 1$. Moreover, again by Proposition 5.5, we know that $(0,1+\rho) \in \operatorname{Cone}(\mathcal{P})$, and so $(0,l) \in \operatorname{Supp}(\ell_{\rho,\sigma}(P))$ for some $l \in \mathbb{N}$. The proof of item (3) is similar. □

We will now see how the different automorphisms of W affect the shape of P (and hence also of Q). We will consider only two types of automorphisms (which are shown in [8] to generate all automorphisms of W). They are the maps

$$\Phi_{n,\lambda}\colon W \to W \quad \text{and} \quad \Phi'_{n,\lambda}\colon W \to W,$$

defined for all $\Lambda \in K$ and $n \in \mathbb{N}$, by

$$\Phi_{n,\lambda}(Y) = Y \quad \text{and} \quad \Phi_{n,\lambda}(X) = X + \lambda Y^n$$

and

$$\Phi'_{n,\lambda}(X) = X \quad \text{and} \quad \Phi'_{n,\lambda}(Y) = Y + \lambda X^n,$$

respectively.

Corollary 5.8. *Let (P,Q) be an irreducible pair and let $(\rho,\sigma) \in \operatorname{Val}(P)$. Take C, m and n be as in Proposition 3.10. We have:*

1. *If $\rho > \sigma > 0$, then $\sigma = 1$ and there exists $\lambda \in K$ such that the automorphism $\varphi := \Phi_{\rho,-\lambda}$ of W, satisfies*
   - (a) *$\ell_{\rho,1}(\varphi(P)) = \mu_P x^{tm} y^{sm}$ and $\ell_{\rho,1}(\varphi(Q)) = \mu_Q x^{tn} y^{sn}$, where $s \in \mathbb{N}_0$, $t = \deg(f_{C,\rho,1}) > 0$ and $\mu_P, \mu_Q \in K^{\times}$.*
   - (b) *$\ell_{\rho_1,\sigma_1}(\varphi(P)) = \ell_{\rho_1,\sigma_1}(P)$ and $\ell_{\rho_1,\sigma_1}(\varphi(Q)) = \ell_{\rho_1,\sigma_1}(Q)$ for all (ρ_1,σ_1) such that $(1,-1) < (\rho_1,\sigma_1) < (\rho,1)$.*
2. *If $\sigma > \rho > 0$, then $\rho = 1$ and there exists $\lambda \in K$ such that the automorphism $\varphi := \Phi'_{\sigma,-\lambda}$ of W, satisfies*
   - (a) *$\ell_{1,\sigma}(\varphi(P)) = \mu_P x^{tn} y^{sn}$ and $\ell_{1,\sigma}(\varphi(Q)) = \mu_Q x^{tm} y^{sm}$, where $t \in \mathbb{N}_0$, $s = \deg(f_{C,1,\sigma}) > 0$ and $\mu_P, \mu_Q \in K^{\times}$.*
   - (b) *$\ell_{\rho_1,\sigma_1}(\varphi(P)) = \ell_{\rho_1,\sigma_1}(P)$ and $\ell_{\rho_1,\sigma_1}(\varphi(Q)) = \ell_{\rho_1,\sigma_1}(Q)$ for all (ρ_1,σ_1) such that $(1,\sigma) < (\rho_1,\sigma_1) < (-1,1)$.*

Proof. We prove item (1) and leave the task to prove item (2) to the reader. Let F be as in item (2) of Theorem 4.5. By its very definition we know that $f_{F,\rho,1}(0) \neq 0$. Hence, by Lemma 5.6 and Proposition 5.7,

$$\sigma = 1 \qquad \text{and} \qquad f_{F,\rho,1} = \mu_F(\lambda x + 1) \quad \text{with } \mu_F, \lambda \in K^{\times}.$$

Thus, by Proposition 4.6, there exist $\mu_P \in K^{\times}$ and $j \in \mathbb{N}$ such that

$$f_{P,\rho,1} = \mu_P(\lambda x + 1)^j.$$

Consequently, there exist $k, u \in \mathbb{N}_0$, such that

$$\ell_{\rho,1}(P) = x^k y^u f_{P,\rho,1}(x^{-1}y^{\rho}) = \mu_P x^k y^u (\lambda x^{-1} y^{\rho} + 1)^j = \mu_P x^{k-j} y^u (\lambda y^{\rho} + x)^j.$$

Note that $k \ge j$, since $\ell_{\rho,1}(P)$ is a polynomial. Since, by item (2) of Proposition 5.7, there exists $l \in \mathbb{N}$ such that

$$(0,l) \in \operatorname{Supp}(\ell_{\rho,1}(P)),$$

we have $k = j$. Hence, by item (1) of Proposition 5.2,

$$\ell_{\rho,1}(\varphi(P)) = \varphi_L(\ell_{\rho,1}(P)) = \mu_P x^j y^u,$$

since $\varphi_L(\lambda y^\rho + x) = x$. We assert that $j = mt$ and $u = ms$, where $s \in \mathbb{N}_0$ and $t = \deg(f_{C,\rho,1}) > 0$. In fact, since, by Proposition 3.10 and 1.9,

$$\ell_{\rho,1}(P) = \lambda_P \ell_{\rho,1}(C)^m, \tag{5.10}$$

we have

$$\ell_{\rho,1}(C) = \mu y^s (\lambda y^\rho + x)^t = \sum_{i=0}^{t} \mu \binom{t}{i} x^{t-i} y^{s+i\rho},$$

where $t = \deg(f_{C,\rho,1}) > 0$, and so, $j = tm$ and $u = sm$ by equality (5.10). Since, $\operatorname{Val}(Q) = \operatorname{Val}(P)$ by Remark 3.9, a similar argument shows that

$$\ell_{\rho,1}(\varphi(Q)) = \mu_Q x^{tn} y^{sn}.$$

Finally, item (b) follows from the additional affirmation in Proposition 5.2. □

Lemma 5.9. *Let $P, Q \in W$. If for some $\mu \in K^\times$ and $r \in \mathbb{N}_0$,*

$$\ell_{1,1}(P) = \begin{cases} \mu(x+\lambda y)^r & \text{for some } \lambda \in K^\times, \\ \mu x^r, \\ \mu y^r, \end{cases}$$

then (P, Q) cannot be an irreducible pair.

Proof. We consider first the case $\ell_{1,1}(P) = \mu y^r$. Assume that (P, Q) is an irreducible pair, which clearly implies that $r > 0$. By Propositions 3.10 and 1.9 there exist $\lambda_P, \lambda_Q \in K^\times$, $C \in W$ and $n, m \in \mathbb{N}$ such that

$$\ell_{1,1}(P) = \lambda_P \ell_{1,1}(C)^m \quad \text{and} \quad \ell_{1,1}(Q) = \lambda_Q \ell_{1,1}(C)^n.$$

Hence

$$\ell_{1,1}(Q) = \lambda_Q \ell_{1,1}(C)^n = \frac{\lambda_Q}{\lambda_P} \ell_{1,1}(P)^{\frac{n}{m}} = \frac{\mu\lambda_Q}{\lambda_P} y^{\frac{rn}{m}} = \mu' y^s, \tag{5.11}$$

where $\mu' := \frac{\mu\lambda_Q}{\lambda_P}$ and $s := \frac{rn}{m}$ (note that $\mu' \in K^\times$ and $s \in \mathbb{N}$). Now, by Remark 3.9 we know that $\operatorname{Val}(P) = \operatorname{Val}(Q)$. Let

$$(\rho,\sigma) := \max\{(\rho',\sigma') \in \overline{\operatorname{Val}}(P) \text{ such that } (\rho',\sigma') < (1,1)\}.$$

By definition $(\rho,\sigma) < (1,1)$ means that $\rho > \sigma$. We claim that $\sigma > 0$ (in particular $\rho + \sigma > 0$ and so $(\rho,\sigma) \in \operatorname{Val}(P)$). Let

$$(\rho'',\sigma'') = \min\{(\rho',\sigma') \in \overline{\operatorname{Val}}(P) \text{ such that } (\rho,\sigma) < (\rho',\sigma')\}.$$

Since $(1,1) \notin \operatorname{Val}(P)$, necessarily $(\rho,\sigma) < (1,1) < (\rho'',\sigma'')$. Hence, by Proposition 2.8,

$$\{\operatorname{en}_{\rho,\sigma}(P)\} = \operatorname{Supp}(\ell_{1,1}(P)) = \{(0,r)\} \tag{5.12}$$

and

$$\{\operatorname{en}_{\rho,\sigma}(Q)\} = \operatorname{Supp}(\ell_{1,1}(Q)) = \{(0,s)\}. \tag{5.13}$$

Consequently, by Proposition 3.6,

$$r\sigma = v_{\rho,\sigma}(0,r) = v_{\rho,\sigma}(P) > 0,$$

which implies that $\sigma > 0$. Now, it follows from item (1) of Corollary 5.8 that $\sigma = 1$ and there exists $\varphi := \Phi_{\rho,-\lambda}$, such that

$$\ell_{\rho,1}(\varphi(P)) = \mu_P x^{tn} y^{ln} \quad \text{where } t \in \mathbb{N},\ l \in \mathbb{N}_0,\ n \in \mathbb{N} \text{ and } \mu_P \in K^{\times}. \tag{5.14}$$

Consequently,

$$\bigl(0, v_{\rho,1}(\varphi(P))\bigr) \notin \operatorname{Supp}(\varphi(P)). \tag{5.15}$$

In fact, if $(0, v_{\rho,1}(\varphi(P))) \in \operatorname{Supp}(\varphi(P))$, then from

$$v_{\rho,1}\bigl(0, v_{\rho,1}(\varphi(P))\bigr) = v_{\rho,1}(\varphi(P)),$$

follows

$$\bigl(0, v_{\rho,1}(\varphi(P))\bigr) \in \operatorname{Supp}\bigl(\ell_{\rho,1}(\varphi(P))\bigr),$$

which contradicts (5.14). On the other hand, by Proposition 5.2 and equalities (5.11), (5.12) and (5.13), we have

$$v_{\rho,1}(\varphi(P)) = v_{\rho,1}(P) = v_{\rho,1}(0,r) = r = v_{1,1}(\mu y^r) = v_{1,1}(P) \tag{5.16}$$

and

$$v_{\rho,1}(\varphi(Q)) = v_{\rho,1}(Q) = v_{\rho,1}(0,s) = s = v_{1,1}(\mu' y^s) = v_{1,1}(Q) \tag{5.17}$$

Moreover, for each $(i,j) \in \operatorname{Supp}(\varphi(P))$,

$$v_{1,1}(X^iY^j) = i + j \le \rho i + j = v_{\rho,1}(X^iY^j) \le v_{\rho,1}(\varphi(P)), \tag{5.18}$$

and, for each $(i,j) \in \operatorname{Supp}(\varphi(Q))$,

$$v_{1,1}(X^iY^j) = i + j \le \rho i + j = v_{\rho,1}(X^iY^j) \le v_{\rho,1}(\varphi(Q)). \tag{5.19}$$

Since $\rho > 1$, the equality in (5.18) is only possible if $i = 0$ and $j = v_{\rho,1}(\varphi(P))$. Thus, by (5.15), (5.16), (5.17), (5.18) and (5.19),

$$v_{1,1}(\varphi(P)) < v_{1,1}(P) \quad \text{and} \quad v_{1,1}(\varphi(Q)) < v_{1,1}(Q),$$

which shows that (P,Q) is not irreducible. The proof in the case $\ell_{1,1}(P) = \mu x^r$ is similar. Finally, we consider the case $\ell_{1,1}(P) = \mu(x + \lambda y)^r$ with $\lambda \in K^{\times}$. Assume that (P,Q) is an irreducible pair and set $\varphi := \Phi_{1,-\lambda}$. By item (1) of Proposition 5.2,

$$\mu x^r = \varphi_L(\mu(x+\lambda y)^r) = \varphi_L(\ell_{1,1}(P)) = \ell_{1,1}(\varphi(P)), \tag{5.20}$$

$$v_{1,1}\bigl(\varphi(P)\bigr) = v_{1,1}(P) \quad \text{and} \quad v_{1,1}\bigl(\varphi(Q)\bigr) = v_{1,1}(Q). \tag{5.21}$$

Now, from (5.21) it follows easily that $(\varphi(P), \varphi(Q))$ is an irreducible pair, which by (5.20) contradicts the previous case, finishing the proof. □

Lemma 5.10. *Assume that* (P,Q) *is irreducible. If*

$$\ell_{1,1}(P) = \mu x^r y^s \quad \textit{with } \mu \in K^{\times} \textit{ and } r, s \in \mathbb{N},$$

then $\operatorname{Supp}(P) \subseteq \{(i,j) : i \le r \textit{ and } j \le s\}$.

Proof. Let $(i,j) \in \operatorname{Supp}(P)$. We prove that $j \le s$ and leave the proof that $i \le r$, which is similar, to the reader. We take

$$(\rho,\sigma) := \min\{(\rho',\sigma') \in \overline{\operatorname{Val}}(P) \text{ such that } (\rho',\sigma') > (1,1)\}.$$

By definition this implies that $\rho < \sigma$. Since $(1,1) \notin \operatorname{Val}(P)$,

$$(\rho'',\sigma'') < (1,1) < (\rho,\sigma),$$

where

$$(\rho'',\sigma'') = \max\{(\rho',\sigma') \in \overline{\operatorname{Val}}(P) \text{ such that } (\rho',\sigma') < (\rho,\sigma)\}.$$

Hence, by Proposition 2.8,

$$\{\operatorname{st}_{\rho,\sigma}(P)\} = \operatorname{Supp}(\ell_{1,1}(P)) = \{(r,s)\}. \tag{5.22}$$

We claim that $(\rho,\sigma)\ge(0,1)$. Otherwise $0<\rho$, which by item (3) of Proposition 5.7, implies $\rho=1$ and

$$(l,0)\in\operatorname{Supp}(\ell_{1,\sigma}(P))\quad\text{for some } l\in\mathbb{N}. \tag{5.23}$$

Now by (5.22) and (5.23), we have

$$r+\sigma s=v_{1,\sigma}(P)=v_{1,\sigma}(l,0)=l,$$

and so

$$v_{1,1}(P)\ge v_{1,1}(l,0)=l=r+\sigma s>r+s$$

since $\sigma>1$. But this contradicts $v_{1,1}(P)=r+s$ and so proves the claim. Then

$$(\rho'',\sigma'')<(1,1)<(0,1)\le(\rho,\sigma),$$

and consequently, by Proposition 2.8,

$$\{\mathrm{st}_{0,1}(P)\}=\operatorname{Supp}(\ell_{1,1}(P))=\{(r,s)\}.$$

Hence,

$$s=v_{0,1}(r,s)=v_{0,1}(P)\ge v_{0,1}(i,j)=j,$$

as desired. □

Theorem 5.11. *Let (P,Q) be an irreducible pair. Then there exist $\mu_P,\mu_Q\in K^\times$, $a,b,m,n\in\mathbb{N}$ and $\varphi\in\operatorname{Aut}(W)$ such that $m,n>1$, $\gcd(m,n)=1$, $1\le a<b$ and*

$$\ell_{1,1}(\varphi(P))=\mu_P x^{am}y^{bm},\qquad \ell_{1,1}(\varphi(Q))=\mu_Q x^{an}y^{bn},$$
$$v_{1,1}(\varphi(P))=v_{1,1}(P),\qquad v_{1,1}(\varphi(Q))=v_{1,1}(Q).$$

Furthermore $\varphi(P)$ and $\varphi(Q)$ are subrectangular and the pair $(\varphi(P),\varphi(Q))$ is irreducible.

Proof. We claim that there exist $\mu\in K^\times$, $r,s\in\mathbb{N}$ and $\varphi\in\operatorname{Aut}(W)$ such that

$$\ell_{1,1}(\varphi(P))=\mu x^r y^s,\quad v_{1,1}(\varphi(P))=v_{1,1}(P)\quad\text{and}\quad v_{1,1}(\varphi(Q))=v_{1,1}(Q).$$

By item (2) of Theorem 4.5 there exists a $(1,1)$-homogeneous element $F\in W$, such that

$$[P,F]_{1,1}=\ell_{1,1}(P)\qquad\text{and}\qquad[Q,\zeta^{-1}F]_{1,1}=\ell_{1,1}(Q),$$

where $\zeta=v_{1,1}(P)/v_{1,1}(Q)$. By item (1) of Proposition 5.7, we know that

$$\operatorname{Supp}(F)\subseteq\{(2,0),(1,1),(0,2)\}.$$

So we have the following cases:

1. $\operatorname{Supp}(F)=\{(2,0)\}$,
2. $\operatorname{Supp}(F)=\{(0,2)\}$,
3. $\operatorname{Supp}(F)=\{(1,1)\}$,
4. $\operatorname{Supp}(F)=\{(1,1),(0,2)\}$,
5. $\operatorname{Supp}(F)=\{(2,0),(1,1)\}$,
6. $\operatorname{Supp}(F)=\{(2,0),(0,2)\}$,
7. $\operatorname{Supp}(F)=\{(2,0),(1,1),(0,2)\}$.

If F is a monomial, then, by Lemma 5.6, $\operatorname{Supp}(F)=\{(1,1)\}$ and $(1,1)\notin\operatorname{Val}(P)$. Consequently, the first two cases are impossible and, in the third one,

$$\ell_{1,1}(P)=\mu x^r y^s\quad\text{with }\mu\in K^\times\text{ and }r,s\in\mathbb{N}_0.$$

Note that by Lemma 5.9, necessarily $r,s>0$, and so we can take $\varphi=\mathrm{id}$. By its very definition, in case (4)

$$f_{F,1,1}=\mu_F(\lambda x+1)\quad\text{with }\mu_F,\lambda\in K^\times.$$

Hence, by Proposition 4.6, there exist $\mu,\bar\mu\in K^\times$ and $r,\bar r\in\mathbb{N}$ such that

$$f_{P,1,1}=\mu(\lambda x+1)^r\qquad\text{and}\qquad f_{Q,1,1}=\bar\mu(\lambda x+1)^{\bar r}.$$

Consequently, there exist $k, \bar{k}, s, \bar{s} \in \mathbb{N}_0$, such that

$$\ell_{1,1}(P) = x^k y^s f_{P,1,1}(x^{-1}y) = \mu x^k y^s (\lambda x^{-1}y + 1)^r = \mu x^{k-r} y^s (\lambda y + x)^r$$

and

$$\ell_{1,1}(Q) = x^{\bar{k}} y^{\bar{s}} f_{Q,1,1}(x^{-1}y) = \bar{\mu} x^{\bar{k}} y^{\bar{s}} (\lambda x^{-1}y + 1)^{\bar{r}} = \mu x^{\bar{k}-\bar{r}} y^{\bar{s}} (\lambda y + x)^{\bar{r}}.$$

Note that $k \ge r$ and $\bar{k} \ge \bar{r}$, since $\ell_{1,1}(P)$ and $\ell_{1,1}(Q)$ are polynomials. By item (1) of Proposition 5.7, there exists $l \in \mathbb{N}$ such that $(0, l) \in \operatorname{Supp}(\ell_{1,1}(P))$. So, $k = r$ and thus $v_{1,1}(P) = r + s$. Similarly $\bar{k} = \bar{r}$ and $v_{1,1}(P) = \bar{r} + \bar{s}$. Hence

$$\ell_{1,1}(P) = \mu(\lambda y + x)^r y^s \qquad \text{and} \qquad \ell_{1,1}(Q) = \bar{\mu}(\lambda y + x)^{\bar{r}} y^{\bar{s}}.$$

On the other hand, by Lemma 5.9, necessarily $s > 0$. Set $\varphi := \Phi_{1,-\lambda}$. By Proposition 5.2,

$$\mu x^r y^s = \varphi_L(\mu(x + \lambda y)^r y^s) = \varphi_L(\ell_{1,1}(P)) = \ell_{1,1}(\varphi(P))$$

and

$$\bar{\mu} x^{\bar{r}} y^{\bar{s}} = \varphi_L(\bar{\mu}(x + \lambda y)^{\bar{r}} y^{\bar{s}}) = \varphi_L(\ell_{1,1}(Q)) = \ell_{1,1}(\varphi(Q)).$$

Consequently

$$v_{1,1}\bigl(\varphi(P)\bigr) = r + s = v_{1,1}(P) \quad \text{and} \quad v_{1,1}\bigl(\varphi(Q)\bigr) = \bar{r} + \bar{s} = v_{1,1}(Q).$$

The case (5) is similar. Finally, we consider the cases (6) and (7). By its very definition $\deg\bigl(f_{F,1,1}\bigr) = 2$. Hence, by Proposition 4.6, there exist $\mu_F, \mu, \bar{\mu}, \lambda_1, \lambda_2 \in K^\times$ and $r, s, \bar{r}, \bar{s} \in \mathbb{N}_0$, with $\lambda_1 \ne \lambda_2$, $r + s > 0$ and $\bar{r} + \bar{s} > 0$, such that

$$f_{F,1,1} = \mu_F(\lambda_1 x + 1)(\lambda_2 x + 1),$$
$$f_{P,1,1} = \mu(\lambda_1 x + 1)^r (\lambda_2 x + 1)^s$$

and

$$f_{Q,1,1} = \bar{\mu}(\lambda_1 x + 1)^{\bar{r}} (\lambda_2 x + 1)^{\bar{s}}.$$

So, there exist $u, v, \bar{u}, \bar{v} \in \mathbb{N}_0$, such that

$$\ell_{1,1}(P) = \mu x^u y^v (\lambda_1 x^{-1} y + 1)^r (\lambda_2 x^{-1} y + 1)^s = \mu x^{u-r-s} y^v (\lambda_1 y + x)^r (\lambda_2 y + x)^s.$$

and

$$\ell_{1,1}(Q) = \bar{\mu} x^{\bar{u}} y^{\bar{v}} (\lambda_1 x^{-1} y + 1)^{\bar{r}} (\lambda_2 x^{-1} y + 1)^{\bar{s}} = \mu x^{\bar{u}-\bar{r}-\bar{s}} y^{\bar{v}} (\lambda_1 y + x)^{\bar{r}} (\lambda_2 y + x)^{\bar{s}}.$$

Note that $u \ge r + s$ and $\bar{u} \ge \bar{r} + \bar{s}$, since $\ell_{1,1}(P)$ and $\ell_{1,1}(Q)$ are polynomials. By item (1) of Proposition 5.7, there exists $l \in \mathbb{N}$ such that $(l, 0), (0, l) \in \operatorname{Supp}(\ell_{1,1}(P))$. So $u = r + s$ and $v = 0$, and thus $v_{1,1}(P) = r + s$. Similarly $\bar{u} = \bar{r} + \bar{s}$, $\bar{v} = 0$ and $v_{1,1}(Q) = \bar{r} + \bar{s}$. Hence

$$\ell_{1,1}(P) = \mu(\lambda_1 y + x)^r (\lambda_2 y + x)^s \quad \text{and} \quad \ell_{1,1}(Q) = \bar{\mu}(\lambda_1 y + x)^{\bar{r}} (\lambda_2 y + x)^{\bar{s}}.$$

Furthermore, from Lemma 5.9 it follows that $r, s, \bar{r}, \bar{s} \in \mathbb{N}$. Set $\psi := \Phi_{1,-\lambda_1}$. By item (1) of Proposition 5.2,

$$\ell_{1,1}(\psi(P)) = \psi_L(\ell_{1,1}(P)) = \mu x^r (x + (\lambda_2 - \lambda_1) y)^s$$

and

$$\ell_{1,1}(\psi(Q)) = \psi_L(\ell_{1,1}(Q)) = \bar{\mu} x^{\bar{r}} (x + (\lambda_2 - \lambda_1) y)^{\bar{s}}.$$

Consequently,

$$v_{1,1}(\psi(P)) = r + s = v_{1,1}(P) \quad \text{and} \quad v_{1,1}(\psi(Q)) = \bar{r} + \bar{s} = v_{1,1}(Q).$$

Now set $P' := \psi(P)$, $Q' := \psi(Q)$, $\lambda := 1/(\lambda_1 - \lambda_2)$ and $\psi' := \Phi'_{1,\lambda}$. By Proposition 5.2,

$$\ell_{1,1}(\psi'(P')) = \psi'_L(\ell_{1,1}(P')) = \mu(\lambda_2 - \lambda_1)^r x^{t_1} y^s$$

and

$$\ell_{1,1}(\psi'(Q')) = \psi'_L(\ell_{1,1}(Q')) = \bar{\mu}(\lambda_2 - \lambda_1)^{\bar{s}} x^{\bar{r}} y^{\bar{s}}.$$

Consequently,

$$v_{1,1}(\psi'(P')) = v_{1,1}(P') = v_{1,1}(P) \quad \text{and} \quad v_{1,1}(\psi'(Q')) = v_{1,1}(Q') = v_{1,1}(Q).$$

So, we can take the automorphism $\varphi := \psi' \circ \psi$, finishing the proof of the claim. Now the pair $(\varphi(P), \varphi(Q))$ is irreducible, since

$$v_{1,1}(\varphi(P)) = v_{1,1}(P), \quad v_{1,1}(\varphi(Q)) = v_{1,1}(Q)$$

and (P, Q) is. Hence, by items (2) and (3) of Proposition 3.10 and item (2) of Proposition 1.9 there exist a $(1,1)$-homogeneous element $C \in W$, $\lambda_P, \lambda_Q \in K^{\times}$ and $n, m \in \mathbb{N}$, such that

$$\mu x^r y^s = \ell_{1,1}(\varphi(P)) = \lambda_P \ell_{1,1}(C)^m \qquad \text{and} \qquad \ell_{1,1}(\varphi(Q)) = \lambda_Q \ell_{1,1}(C)^n. \tag{5.24}$$

Let $d := \gcd(m, n)$. Replacing C by C^d, we can assume that $\gcd(m, n) = 1$. Note that, by item (4) of Proposition 3.10, we know that $m, n > 1$. From the first equality in (5.24) it follows that there exists $a, b \in \mathbb{N}$ such that

$$r = ma, \quad s = mb \quad \text{and} \quad \ell_{1,1}(C) = \mu \lambda_P^{-1} x^a y^b.$$

So, by (5.24),

$$\ell_{1,1}(\varphi(P)) = \mu x^{ma} y^{mb} \quad \text{and} \quad \ell_{1,1}(\varphi(Q)) = \lambda_Q \mu^n \lambda_P^{-n} x^{na} y^{nb}.$$

Consequently, by Lemma 5.10, $\varphi(P)$ and $\varphi(Q)$ are subrectangular. Furthermore, by item (1) of Theorem 4.5, we have $a \neq b$. Finally, thanks to the existence of an automorphism of W that takes X to Y and Y to $-X$, we can obtain $a < b$. ☐

Corollary 5.12. *If the Dixmier conjecture is false, then there exist $P, Q \in W$ such that $[Q, P] = 1$ and the supports of P and Q are subrectangular but not subsquare.*

Proof. It follows from Theorems 3.3 and 5.11. ☐

6 Computing lower bounds

In this section we assume that K is algebraically closed. Our purpose in this section is to determine a lower bound for the value

$$B := \min\{\gcd(v_{1,1}(P), v_{1,1}(Q)), \text{ where } (P, Q) \text{ is an irreducible pair}\}.$$

More precisely, we will prove that $B > 8$.

Lemma 6.1. *Let $P \in W$ with $v_{1,-1}(\mathrm{en}_{1,0}(P)) < 0$. Suppose there exists $F \in K[Y]$ satisfying*

$$[P, XF(Y)]_{1,0} = \ell_{1,0}(P). \tag{6.1}$$

Then there is a $\lambda \in K$ such that the automorphism φ of W, defined by

$$\varphi(X) := X \qquad \text{and} \qquad \varphi(Y) := Y + \lambda,$$

fulfills $v_{1,-1}(\mathrm{st}_{1,0}(\varphi(P))) \leq 0$.

Proof. Let $(a, b) := \mathrm{en}_{1,0}(P)$. By definition there exists $f \in K[y]$, with $\deg(f) = b$, such that $\ell_{1,0}(P) = x^a f(y)$. Moreover, $b > a$ since

$$a - b = v_{1,-1}(\mathrm{en}_{1,0}(P)) < 0.$$

We claim that there is a linear factor $y - \lambda$ of f, whose multiplicity m_λ fulfills $m_\lambda \geq a$. This is clear if $a = 0$. So, we can assume that $a > 0$. A direct computation using Proposition 1.18 shows that

$$[P, XF(Y)]_{1,0} = [X^a f(Y), XF(Y)]_{1,0} = x^a (f'(y)F(y) - af(y)F'(y)),$$

and so, hypothesis (6.1) reads

$$f'F - afF' = f.$$

From this it follows that any linear factor of f divides F. Hence,

$$\#\,\text{factors}(f) \le \deg(F), \tag{6.2}$$

where $\#\,\text{factors}(f)$ denotes the number of different linear factors of f. Note that

$$\text{en}_{1,0}(XF(Y)) = (1,M),$$

where $M := \deg(F)$. By item (2) of Proposition 1.17, we know that

$$(1,M) = (1,1) \quad \text{or} \quad (1,M) \sim \text{en}_{1,0}(P) = (a,b).$$

This yields $M = 1$ or $M = b/a > 1$. Consequently, by (6.2),

$$\#\,\text{factors}(f) \le b/a.$$

Hence, there exists a linear factor $y - \lambda$ of f, whose multiplicity m_λ fulfills

$$m_\lambda \ge \frac{\deg(f)}{\#\,\text{factors}(f)} \ge \frac{b}{b/a} = a,$$

which finishes the proof of the claim.

Write $f(y) := (y-\lambda)^{m_\lambda}\overline{f}(y)$. By Proposition 5.2, the automorphism φ of W, defined by $\varphi(X) := X$ and $\varphi(Y) := Y + \lambda$, satisfies

$$\ell_{1,0}(\varphi(P)) = \varphi_L(\ell_{1,0}(P)) = x^a y^{m_\lambda}\overline{f}(y+\lambda).$$

Thus $\text{st}_{1,0}(\varphi(P)) = (a, m_\lambda)$, and so,

$$v_{1,-1}(\text{st}_{1,0}(\varphi(P))) = a - m_\lambda \le 0,$$

as desired. □

Proposition 6.2. *For any irreducible pair (P,Q), there exist $m,n \in \mathbb{N}$ and an automorphism ψ of W, such that $m,n > 1$, $\gcd(m,n) = 1$,*

1. *$(P_0,Q_0) := (\psi(P),\psi(Q))$ is an irreducible,*
2. *$v_{1,1}(P_0) = v_{1,1}(P)$ and $v_{1,1}(Q_0) = v_{1,1}(Q)$,*
3. *P_0 and Q_0 are subrectangular and*
   $$\text{Supp}(\ell_{1,1}(P_0)) = (ma,mb) \quad \textit{and} \quad \text{Supp}(\ell_{1,1}(Q_0)) = (na,nb),$$
   for some $a,b \in \mathbb{N}$ with $a < b$,
4. *$v_{1,-1}(\text{en}_{1,0}(P_0)) < 0$ and $v_{1,-1}(\text{en}_{1,0}(Q_0)) < 0$,*
5. *$v_{1,-1}(\text{st}_{1,0}(P_0)) < 0$ and $v_{1,-1}(\text{st}_{1,0}(Q_0)) < 0$.*

Moreover, there exists $(\rho,\sigma) := \text{Pred}_{1,0}(P_0)$, and

- (a) *$\sigma < 0$,*
- (b) *$[Q_0,P_0] = 1$,*
- (c) *$(\rho,\sigma) \in \text{Val}(P_0) \cap \text{Val}(Q_0)$,*
- (d) *$v_{\rho,\sigma}(P_0) > 0$ and $v_{\rho,\sigma}(Q_0) > 0$,*
- (e) *$[P_0,Q_0]_{\rho,\sigma} = 0$,*
- (f) *$\frac{v_{\rho,\sigma}(P_0)}{v_{\rho,\sigma}(Q_0)} = \frac{m}{n}$,*
- (g) *$v_{1,-1}\big(\text{en}_{\rho,\sigma}(P_0)\big) < 0$ and $v_{1,-1}\big(\text{en}_{\rho,\sigma}(Q_0)\big) < 0$.*

Proof. By Theorem 5.11 there exist $\mu_P,\mu_Q \in K^\times$, $a,b,m,n \in \mathbb{N}$ and $\psi_1 \in \text{Aut}(W)$ such that $m,n > 1$, $\gcd(m,n) = 1$, $1 \le a < b$ and

$$\ell_{1,1}(\psi_1(P)) = \mu_P x^{am} y^{bm}, \qquad \ell_{1,1}(\psi_1(Q)) = \mu_Q x^{an} y^{bn}, \tag{6.3}$$

$$v_{1,1}(\psi_1(P)) = v_{1,1}(P), \qquad v_{1,1}(\psi_1(Q)) = v_{1,1}(Q). \tag{6.4}$$

Furthermore

- the pair $(\psi_1(P),\psi_1(Q))$ is irreducible,

- $\psi_1(P)$ and $\psi_1(Q)$ are subrectangular.

Set

$$P_1 := \psi_1(P) \qquad \text{and} \qquad Q_1 := \psi_1(Q).$$

Since P_1 is subrectangular and $\operatorname{Supp}(\ell_{1,1}(P_1)) = \{(am,bm)\}$, we have

$$\{(am,bm)\} \subseteq \operatorname{Supp}(\ell_{1,0}(P_1)) \subseteq \{(am,j) : 0 \le j \le bm\}.$$

Hence,

$$\operatorname{en}_{1,0}(P_1)) = (am,bm), \quad \text{and similarly} \quad \operatorname{en}_{1,0}(Q_1)) = (an,bn).$$

Consequently, by the first equality,

$$v_{1,-1}(\operatorname{en}_{1,0}(P_1)) = v_{1,-1}(am,bm) = m(a-b) < 0.$$

On the other hand, by item (3) of Theorem 4.5 there exists a $(1,0)$-homogeneous element F_1 such that

$$[P,F_1]_{1,0} = \ell_{1,0}(P) \quad \text{and} \quad v_{1,0}(F_1) = 1.$$

This implies that $F_1 = XF(Y)$ with $F \in K[Y]$. Hence, by Lemma 6.1, there exists $\lambda \in K$ such that the automorphism φ of W, defined by

$$\varphi(X) := X \qquad \text{and} \qquad \varphi(Y) := Y + \lambda,$$

fulfills

$$v_{1,-1}(\operatorname{st}_{1,0}(\varphi(P_1))) \le 0. \tag{6.5}$$

Moreover, since by Proposition 5.2,

$$\ell_{1,1}(\varphi(P_1)) = \ell_{1,1}(P_1) = \mu_P x^{am} y^{bm} \tag{6.6}$$

and

$$\ell_{1,1}(\varphi(Q_1)) = \ell_{1,1}(Q_1) = \mu_Q x^{an} y^{bn}, \tag{6.7}$$

the pair $(P_0,Q_0) := (\varphi(P_1),\varphi(Q_1))$ is irreducible. Consequently, by item (2) of Theorem 4.5, inequality (6.5) and Remark 3.11,

$$v_{1,-1}(\operatorname{st}_{1,0}(P_0)) < 0 \quad \text{and} \quad v_{1,-1}(\operatorname{st}_{1,0}(Q_0)) < 0, \tag{6.8}$$

while, by (6.6), Lemma 5.10 and Remark 3.9, the elements P_0 and Q_0 are subrectangular. Now, arguing as above for (P_1,Q_1), we obtain that

$$\operatorname{en}_{1,0}(P_0)) = (am,bm) \quad \text{and} \quad \operatorname{en}_{1,0}(Q_0)) = (an,bn),$$

and so

$$v_{1,-1}(\operatorname{en}_{1,0}(P_0)) < 0 \quad \text{and} \quad v_{1,-1}(\operatorname{en}_{1,0}(Q_0)) < 0.$$

Set $\psi := \varphi \circ \psi_1$. By (6.4), (6.6) and (6.7),

$$v_{1,1}(P_0) = v_{1,1}(P) \quad \text{and} \quad v_{1,1}(Q_0) = v_{1,1}(Q)$$

This concludes the proof of items (1), (2), (3), (4) and (5). As above of Lemma 2.4, let

$$\operatorname{Valinf}_{1,0}(P_0) := \left\{\operatorname{val}\left((i,j) - \operatorname{st}\right) : (i,j) \in \operatorname{Supp}(P_0) \text{ and } v_{1,-1}\left(i,j\right) > v_{1,-1}(\operatorname{st})\right\},$$

where $\operatorname{st} := \operatorname{st}_{1,0}(P_0)$. Note that $\operatorname{Valinf}_{1,0}(P) \ne \emptyset$. In fact, otherwise, by item (5),

$$v_{1,-1}(P_0) = v_{1,-1}(\operatorname{st}_{1,0}(P_0)) < 0,$$

which contradicts Proposition 3.6. Let

$$(\rho,\sigma) = \operatorname{Pred}_{1,0}(P_0) := \max \operatorname{Valinf}_{1,0}(P_0).$$

Item (a) follows from $(\rho,\sigma) < (1,0)$, which is true by item (2) of Lemma 2.4. Item (b) holds, since (P_0,Q_0) are an irreducible pair. Item (c) is true by item (2) of

Lemma 2.7 and Remark 3.9. Items (d) and (e) follow from Propositions 3.6 and 3.7 respectively. Now, by Proposition 3.10 and equalities (6.6) and (6.7),

$$\frac{v_{\rho,\sigma}(P_0)}{v_{\rho,\sigma}(Q_0)}=\frac{v_{1,1}(P_0)}{v_{1,1}(Q_0)}=\frac{m}{n},$$

and so item (f) holds, since $m,n>1$ and $\gcd(m,n)=1$. Finally, item (g) is true by item (2) of Lemma 2.7 and item (5). ☐

Proposition 6.3. *Let (P,Q) be an irreducible pair and $(\rho,\sigma)\in\mathfrak{V}$. The following assertions hold:*

1. *There exist $\lambda_P,\lambda_Q\in K^{\times}$, $m,n\in\mathbb{N}$ and a (ρ,σ)-homogeneous element $R\in L$ such that $m,n>1$, $\gcd(n,m)=1$ and*

   $$\ell_{\rho,\sigma}(P)=\lambda_P R^m \quad \textit{and} \quad \ell_{\rho,\sigma}(Q)=\lambda_Q R^n.$$

   Furthermore

   $$\frac{v_{\rho,\sigma}(P)}{v_{\rho,\sigma}(Q)}=\frac{v_{1,1}(P)}{v_{1,1}(Q)}=\frac{m}{n}.$$

2. *There exists a (ρ,σ)-homogeneous element $F\in W$ such that*

   $$[P,F]_{\rho,\sigma}=\ell_{\rho,\sigma}(P) \quad \textit{and} \quad v_{\rho,\sigma}(F)=\rho+\sigma,$$

   Moreover, if $(\rho,\sigma)\in\operatorname{Val}(P)$ and $\sigma<0$, then F is not a monomial and

   $$\operatorname{en}_{\rho,\sigma}(F)\neq(1,1) \quad \textit{and} \quad \operatorname{en}_{\rho,\sigma}(F)=\mu\operatorname{en}_{\rho,\sigma}(R). \tag{6.9}$$

   where $\mu\in\mathbb{Q}$ and $0<\mu<1$.

Assume that $(\rho,\sigma)\in\operatorname{Val}(P)$ and $\sigma<0$, and write

$$(r,s):=\operatorname{en}_{\rho,\sigma}(R) \quad \textit{and} \quad (\bar r,\bar s):=\operatorname{st}_{\rho,\sigma}(R).$$

The following assertions hold

3. $\gcd(r,s)>1$.
4. $v_{1,0}(\operatorname{en}_{\rho,\sigma}(F))\ge 2$.
5. $r\ge 3$.
6. *If $\bar r\le\bar s$, then $\gcd(\bar r,\bar s)>1$ and $\bar r\ge 3$.*

Proof. By item (2) of Proposition 1.9 and items (2) and (3) of Proposition 3.10, there exist a (ρ,σ)-homogeneous element $C\in W$, $\lambda_P,\lambda_Q\in K^{\times}$ and $n,m\in\mathbb{N}$ such that

$$\ell_{\rho,\sigma}(P)=\lambda_P\ell_{\rho,\sigma}(C^m)=\lambda_P R^m \quad\text{and}\quad \ell_{\rho,\sigma}(Q)=\lambda_Q\ell_{\rho,\sigma}(C^n)=\lambda_Q R^n. \tag{6.10}$$

where $R:=\ell_{\rho,\sigma}(C)$. Let $d:=\gcd(m,n)$. Replacing R by R^d we can assume that $\gcd(m,n)=1$. Note that, by item (4) of Proposition 3.10, we have $m,n>1$. Furthermore, by equality (3.7), item (3) of Proposition 1.9 and (6.10),

$$\frac{m}{n}=\frac{v_{\rho,\sigma}(P)}{v_{\rho,\sigma}(Q)}=\frac{v_{1,1}(P)}{v_{1,1}(Q)}. \tag{6.11}$$

This finishes the proof of item (1). We now prove item (2). By (6.10) and item (5) of Proposition 1.9,

$$\operatorname{en}_{\rho,\sigma}(P)=m\operatorname{en}_{\rho,\sigma}(R) \quad\text{and}\quad \operatorname{en}_{\rho,\sigma}(Q)=n\operatorname{en}_{\rho,\sigma}(R), \tag{6.12}$$

while, by Proposition 3.6

$$v_{\rho,\sigma}(P)>0 \quad\text{and}\quad v_{\rho,\sigma}(Q)>0.$$

Note that from the first inequality above and (6.12) it follows that

$$\operatorname{en}_{\rho,\sigma}(R)\neq(0,0),\quad \operatorname{en}_{\rho,\sigma}(P)\neq(0,0)\quad\text{and}\quad \operatorname{en}_{\rho,\sigma}(Q)\neq(0,0). \tag{6.13}$$

On the other hand, by item (3) of Theorem 4.5, there exists a (ρ,σ)-homogeneous element $F\in W$, such that

$$[F,P]_{\rho,\sigma}=\ell_{\rho,\sigma}(P)\quad\text{and}\quad v_{\rho,\sigma}(F)=\rho+\sigma.$$

Consequently $\mathrm{en}_{\rho,\sigma}(F)\neq(0,0)$. We claim that if $(\rho,\sigma)\in\mathrm{Val}(P)$ and $\sigma<0$, then

$$\mathrm{en}_{\rho,\sigma}(F)\neq(1,1)\quad\text{and}\quad\mathrm{en}_{\rho,\sigma}(F)\sim\mathrm{en}_{\rho,\sigma}(R).$$

In fact, by Lemma 5.6, we know that F is not a monomial, and so $\mathrm{en}_{\rho,\sigma}(F)=(1,1)$ implies that $\mathrm{st}_{\rho,\sigma}(F)=(1+\sigma/\rho,0)$, which is impossible, since from $\sigma<0$ and $\rho+\sigma>0$ it follows that $1+\sigma/\rho\notin\mathbb{Z}$. Hence, by item (2) of Proposition 1.17,

$$\mathrm{en}_{\rho,\sigma}(F)\sim\mathrm{en}_{\rho,\sigma}(P).$$

So, by (6.12) and (6.13), we have $\mathrm{en}_{\rho,\sigma}(F)\sim\mathrm{en}_{\rho,\sigma}(R)$, as desired. By the first inequality in (6.13), this implies that there exists $\mu\in Q$ such that

$$\mathrm{en}_{\rho,\sigma}(F)=\mu\,\mathrm{en}_{\rho,\sigma}(R).$$

In order to end the proof of item (2) it remains to see that $0<\mu<1$. The first inequality follows immediately from the fact that $\mathrm{en}_{\rho,\sigma}(F)\neq(0,0)$. We now are going to prove that $\mu<1$. Let $(\rho',\sigma')\in\mathfrak{V}$ such that

$$(\rho',\sigma')<(\rho'',\sigma'')\quad\text{for all }(\rho'',\sigma'')\in\mathrm{Val}(P).$$

By item (3) of Proposition 2.8 we know that

$$\{w(P)\}=\mathrm{Supp}(\ell_{\rho',\sigma'}(P)).$$

On the other hand, by item (1) there exists $\zeta'\in K^{\times}$ and a (ρ',σ')-homogeneous element $R'\in L$ such that

$$\ell_{\rho',\sigma'}(P)=\zeta' R'^{m}.$$

Hence

$$w(P)=(mr',ms')\quad\text{where }\{(r',s')\}=\mathrm{Supp}(R').$$

Furthermore $r'>s'$, since $v_{1,-1}(P)>0$ by Proposition 3.6. We claim that

$$v_{\rho,\sigma}(r',s')>\rho+\sigma.$$

In fact, if $s'=0$, then $r'\geq 1$ and so

$$v_{\rho,\sigma}(r',s')=\rho r'+\sigma s'=\rho r'\geq\rho>\rho+\sigma,$$

and if $s'\geq 1$, then

$$v_{\rho,\sigma}(r',s')=\rho r'+\sigma s'>\rho s'+\sigma s'=(\rho+\sigma)s'\geq\rho+\sigma.$$

Consequently, by item (3) of Proposition 1.9

$$v_{\rho,\sigma}(R)=\frac{1}{m}v_{\rho,\sigma}(P)\geq\frac{1}{m}v_{\rho,\sigma}(w(P))=v_{\rho,\sigma}(r',s')>\rho+\sigma,$$

and so

$$1>\frac{\rho+\sigma}{v_{\rho,\sigma}(R)}=\frac{v_{\rho,\sigma}(F)}{v_{\rho,\sigma}(R)}=\mu,$$

proving item (2). Now item (3) is clear, since by equality (6.9), the assumption that $\gcd(r,s)=1$ leads to $\mu\in\mathbb{Z}$, which contradicts that $0<\mu<1$. In order to prove item (4) we must show that

$$\mathrm{en}_{\rho,\sigma}(F)=(1,k)\qquad\text{and}\qquad\mathrm{en}_{\rho,\sigma}(F)=(0,k)$$

are impossible. We already know that $\mathrm{en}_{\rho,\sigma}(F)\neq(0,0)$ and $\mathrm{en}_{\rho,\sigma}(F)\neq(1,1)$. If $\mathrm{en}_{\rho,\sigma}(F)=(1,k)$ for some $k\neq 1$, then

$$\rho+k\sigma=v_{\rho,\sigma}(\mathrm{en}_{\rho,\sigma}(F))=v_{\rho,\sigma}(F)=\rho+\sigma,$$

and so $\sigma = 0$, which is a contradiction. Similarly, if $\mathrm{en}_{\rho,\sigma}(F) = (0,k)$ with $k \geq 1$, then

$$k\sigma = v_{\rho,\sigma}(\mathrm{en}_{\rho,\sigma}(F)) = v_{\rho,\sigma}(F) = \rho + \sigma > 0,$$

which contradicts $\sigma < 0$. We now prove item (5). By the very definition of μ,

$$\mathrm{en}_{\rho,\sigma}(F) = (\mu r, \mu s),$$

which, by item (4), implies $r \neq 0$. Hence, again by item (4), we have

$$r > \mu r = v_{1,0}(\mathrm{en}_{\rho,\sigma}(F)) \geq 2.$$

We finally prove item (6). Suppose that $\bar{r} \leq \bar{s}$. As above of Lemma 2.4, let

$$\mathrm{Valinf}_{\rho,\sigma}(P) := \{\mathrm{val}\left((i,j) - \mathrm{st}\right) : (i,j) \in \mathrm{Supp}(P) \text{ and } v_{1,-1}\left(i,j\right) > v_{1,-1}(\mathrm{st})\},$$

where $\mathrm{st} := \mathrm{st}_{\rho,\sigma}(P)$. Since, by item (4) of Proposition 1.9 and Proposition 3.6,

$$v_{1,-1}(\mathrm{st}_{\rho,\sigma}(P)) = m(\bar{r} - \bar{s}) \leq 0 \quad \text{and} \quad v_{1,-1}(P) > 0,$$

we have $\mathrm{Valinf}_{\rho,\sigma}(P) \neq \emptyset$. Let

$$(\bar{\rho}, \bar{\sigma}) := \mathrm{Pred}_{\rho,\sigma}(P) := \max(\mathrm{Valinf}_{\rho,\sigma}(P)).$$

By item (2) of Lemma 2.7 we know that

$$(\bar{\rho}, \bar{\sigma}) \in \mathrm{Val}(P) \quad \text{and} \quad \mathrm{en}_{\bar{\rho},\bar{\sigma}}(P) = \mathrm{st}_{\rho,\sigma}(P).$$

Now, again by item (1), there exists $\bar{\zeta} \in K^{\times}$ and a $(\bar{\rho},\bar{\sigma})$-homogeneous element $\bar{R} \in L \setminus \{0\}$ such that

$$\ell_{\bar{\rho},\bar{\sigma}}(P) = \bar{\zeta}\bar{R}^m.$$

By items (4) and (5) of Proposition 1.9

$$\mathrm{en}_{\bar{\rho},\bar{\sigma}}(\bar{R}) = \frac{1}{m}\,\mathrm{en}_{\bar{\rho},\bar{\sigma}}(P) = \frac{1}{m}\,\mathrm{st}_{\rho,\sigma}(P) = \mathrm{st}_{\rho,\sigma}(R) = (\bar{r},\bar{s}),$$

and so, applying items (3) and (5) with (ρ,σ) replaced by $(\bar{\rho},\bar{\sigma})$, we obtain that $\gcd(\bar{r},\bar{s}) > 1$ and $\bar{r} \geq 3$. ☐

Corollary 6.4. *Let*

$$B := \min\{\gcd(v_{1,1}(P), v_{1,1}(Q)),\ \textit{where } (P,Q) \textit{ is an irreducible pair}\}.$$

Then $B > 8$.

Proof. Let (P,Q) be an irreducible pair such that $B := \gcd(v_{1,1}(P), v_{1,1}(Q))$. Let (P_0,Q_0) and (ρ,σ) be as in Proposition 6.2. Thus, (P_0,Q_0) is an irreducible pair, $\sigma < 0$ and $(\rho,\sigma) \in \mathrm{Val}(P_0) \cap \mathrm{Val}(Q_0)$. Hence, by item (1) of Proposition 6.3, there exists $\lambda_{P_0}, \lambda_{Q_0} \in K^{\times}$, a (ρ,σ)-homogeneous element $R \in L \setminus \{0\}$, and $m,n \in \mathbb{N}$, with $m,n > 1$ and $\gcd(m,n) = 1$, such that

$$\ell_{\rho,\sigma}(P_0) = \lambda_{P_0} R^m, \qquad \ell_{\rho,\sigma}(Q_0) = \lambda_{Q_0} R^n \tag{6.14}$$

and

$$\frac{m}{n} = \frac{v_{\rho,\sigma}(P_0)}{v_{\rho,\sigma}(Q_0)} = \frac{v_{1,1}(P_0)}{v_{1,1}(Q_0)}. \tag{6.15}$$

From the first equality in (6.14) and item (5) of Proposition 1.9, it follows that

$$\frac{1}{m}\,\mathrm{en}_{\rho,\sigma}(P_0) = \mathrm{en}_{\rho,\sigma}(R),$$

while equality (6.15) implies that

$$\gcd(v_{1,1}(P_0), v_{1,1}(Q_0)) = \frac{1}{m} v_{1,1}(P_0).$$

On the other hand, by item (2) of Proposition 6.2,

$$v_{1,1}(P_0) = v_{1,1}(P) \quad \text{and} \quad v_{1,1}(Q_0) = v_{1,1}(Q), \tag{6.16}$$

and so

$$B = \gcd(v_{1,1}(P_0), v_{1,1}(Q_0)) = \frac{1}{m} v_{1,1}(P_0) \geq v_{1,1}\bigl(\mathrm{en}_{\rho,\sigma}(R)\bigr) = r + s,$$

where $(r,s) := \mathrm{en}_{\rho,\sigma}(R)$. Since, by item (g) of Proposition 6.2, we have $r < s$, it follows from items (3) and (5) of Proposition 6.3, that necessarily $r + s \geq 9$. $\square$

References

[1] Harry Appelgate and Hironori Onishi, *The Jacobian conjecture in two variables*, J. Pure Appl. Algebra **37** (1985), no. 3, 215–227, DOI 10.1016/0022-4049(85)90099-4. MR797863 (87b:14005)

[2] Pascal Kossivi Adjamagbo and Arno van den Essen, *A proof of the equivalence of the Dixmier, Jacobian and Poisson conjectures*, Acta Math. Vietnam. **32** (2007), no. 2-3, 205–214. MR2368008 (2009a:14079)

[3] Hyman Bass, Edwin Connell, and David Wright, *The Jacobian conjecture: reduction of degree and formal expansion of the inverse*, Bull. Amer. Math. Soc. (N.S.) **7** (1982), no. 2, 287–330. MR663785 (83k:14028)

[4] Vladimir Bavula, *The Jacobian conjecture*$_{2n}$ *implies the Dixmier problem*$_n$.

[5] V. V. Bavula, *A question of Rentschler and the Dixmier problem*, Ann. of Math. (2) **154** (2001), no. 3, 683–702, DOI 10.2307/3062144. MR1884619 (2003d:16032)

[6] Alexei Belov-Kanel and Maxim Kontsevich, *The Jacobian conjecture is stably equivalent to the Dixmier conjecture*, Mosc. Math. J. **7** (2007), no. 2, 209–218, 349 (English, with English and Russian summaries). MR2337879 (2009f:16041)

[7] Joseph Bernstein and Valery Lunts, *On nonholonomic irreducible D-modules*, Invent. Math. **94** (1988), no. 2, 223–243, DOI 10.1007/BF01394325. MR958832 (90b:58247)

[8] Jacques Dixmier, *Sur les algèbres de Weyl*, Bull. Soc. Math. France **96** (1968), 209–242 (French). MR0242897 (39 #4224)

[9] Jorge A. Guccione, Juan J. Guccione, and Christian Valqui, *On the centralizers in the Weyl algebra*, Proc. Amer. Math. Soc. **140** (2012), no. 4, 1233–1241, DOI 10.1090/S0002-9939-2011-11017-7. MR2869108

[10] Valery Lunts, *Algebraic varieties preserved by generic flows*, Duke Math. J. **58** (1989), no. 3, 531–554, DOI 10.1215/S0012-7094-89-05824-9. MR1016433 (91a:32015)

[11] T. T. Moh, *On the Jacobian conjecture and the configurations of roots*, J. Reine Angew. Math. **340** (1983), 140–212. MR691964 (84m:14018)

[12] Masayoshi Nagata, *Some remarks on the two-dimensional Jacobian conjecture*, Chinese J. Math. **17** (1989), no. 1, 1–7. MR1007872 (90k:14010b)

[13] ———, *Two-dimensional Jacobian conjecture*, Algebra and topology 1988 (Taejǒn, 1988), Korea Inst. Tech., Taejǒn, 1988, pp. 77–98. MR1022249 (90k:14010a)

[14] Yoshifumi Tsuchimoto, *Endomorphisms of Weyl algebra and p-curvatures*, Osaka J. Math. **42** (2005), no. 2, 435–452. MR2147727 (2006g:14101)

[15] Arno van den Essen, *Polynomial automorphisms and the Jacobian conjecture*, Progress in Mathematics, vol. 190, Birkhäuser Verlag, Basel, 2000. MR1790619 (2001j:14082)

[16] Wenhua Zhao, *Images of commuting differential operators of order one with constant leading coefficients*, J. Algebra **324** (2010), no. 2, 231–247, DOI 10.1016/j.jalgebra.2010.04.022. MR2651354 (2011f:13024)

Departamento de Matemática, Facultad de Ciencias Exactas y Naturales-UBA, Pabellón 1-Ciudad Universitaria, Intendente Guiraldes 2160 (C1428EGA) Buenos Aires, Argentina.

Instituto de Investigaciones Matemáticas “Luis A. Santaló”, Facultad de Ciencias Exactas y Naturales-UBA, Pabellón 1-Ciudad Universitaria, Intendente Guiraldes 2160 (C1428EGA) Buenos Aires, Argentina.

E-mail address: `vander@dm.uba.ar`

Departamento de Matemática, Facultad de Ciencias Exactas y Naturales-UBA, Pabellón 1-Ciudad Universitaria, Intendente Guiraldes 2160 (C1428EGA) Buenos Aires, Argentina.

Instituto Argentino de Matemática-CONICET, Savedra 15 3er piso, (C1083ACA) Buenos Aires, Argentina.
E-mail address: `jjgucci@dm.uba.ar`

Pontificia Universidad Católica del Perú - Instituto de Matemática y Ciencias Afines, Sección Matemáticas, PUCP, Av. Universitaria 1801, San Miguel, Lima 32, Perú.

Instituto de Matemática y Ciencias Afines (IMCA) Calle Los Biólogos 245. Urb San César. La Molina, Lima 12, Perú.
E-mail address: `cvalqui@pucp.edu.pe`